

**Table 3:** Body weight of all the studied queens treated with megestrol acetate, before and at the end of treatment, at the first estrus and 4 to 6 months post-treatment.

<b>Cat</b>	<b>Weight before treatment (kg)</b>	<b>Weight at the end of treatment (kg)</b>	<b>Weight at 1st estrus post-treatment (kg)</b>	<b>Weight 4-6 months post-treatment (kg)</b>
<b>P1</b>	2.8	3	NA	NA
<b>P2</b>	3.1	3.3	NA	3.8
<b>P3</b>	3.3	NA	NA	NA
<b>P4</b>	2.8	NA	NA	NA
<b>P5</b>	1.8	NA	NA	NA
<b>P6</b>	3.5	NA	NA	NA
<b>P7</b>	3.1	NA	NA	NA
<b>P8</b>	3.6	NA	NA	NA
<b>P9</b>	2.8	NA	NA	NA
<b>P10</b>	3.1	NA	NA	NA
<b>P11</b>	4.2	4.2	4	NA
<b>P12</b>	3.2	4	4	NA
<b>P13</b>	3.2	3.8	3.7	NA
<b>P14</b>	3.7	5.6	5.5	NA
<b>P15</b>	4.2	4.7	NA	NA
<b>P16</b>	3.4	4.2	4.6	4
<b>P17</b>	2.9	3.1	3.4	3
<b>L1</b>	3.5	4	3.7	3.1
<b>L2</b>	3.7	4.7	4	3.7
<b>L3</b>	4.8	4.9	4.4	4.4
<b>L4</b>	2.3	3	NA	2.5
<b>L5</b>	2.8	3.7	NA	2.9
<b>L6</b>	3.4	3.3	NA	3
<b>L7</b>	4.3	4.7	NA	4.3
<b>L8</b>	2.9	3.8	NA	3.2
<b>L9</b>	2.2	2.5	2.3	2
<b>L10</b>	5.6	5.5	NA	5.4
<b>L11</b>	5.1	5.4	NA	5.1

NA – non-available