

# Calonectria species isolated from *Eucalyptus* plantations and nurseries in South China

JieQiong Li<sup>1,2</sup>, Michael J. Wingfield<sup>2</sup>, QianLi Liu<sup>1</sup>, Irene Barnes<sup>3</sup>, Jolanda Roux<sup>4</sup>, Lorenzo Lombard<sup>5</sup>, Pedro W. Crous<sup>2,5</sup>, and ShuaiFei Chen<sup>1,2</sup>

<sup>1</sup>China Eucalypt Research Centre (CERC), Chinese Academy of Forestry (CAF), ZhanJiang 524022, GuangDong Province, China; corresponding author e-mail: shuaifei.chen@gmail.com

<sup>2</sup>Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, 0028, South Africa

<sup>3</sup>Department of Genetics, FABI, University of Pretoria, Pretoria 0028, South Africa

<sup>4</sup>Department of Plant and Soil Sciences, FABI, University of Pretoria, Pretoria, 0028, South Africa

<sup>5</sup>Westerdijk Fungal Biodiversity Institute, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands

**Abstract:** Diseases caused by species of *Calonectria* (*Ca.*) represent a serious threat to the growth and sustainability of *Eucalyptus* plantations in China. Symptoms caused by these fungi mainly include leaf blight on trees in plantations and rotting of stems and leaves in nurseries. Extensive surveys have recently been conducted where *Calonectria* species were collected in *Eucalyptus* plantations and nurseries in the Fujian, GuangDong, GuangXi, and YunNan Provinces of South China. Additional isolates were baited from soil samples in the Hong Kong Region. The aim of this study was to identify the 115 *Calonectria* isolates obtained using comparisons of DNA sequence data for the  $\beta$ -tubulin (*tub2*), calmodulin (*cmaD*), histone H3 (*his3*) and partial translation elongation factor-1 $\alpha$  (*tef1*) gene regions as well as their morphological features. Seven known species were identified, including *Calonectria arbusta*, *Ca. asiatica*, *Ca. chinensis*, *Ca. eucalypti*, *Ca. hongkongensis*, *Ca. mossambicensis* and *Ca. pentaseptata*. In addition, six novel taxa were collected and are described here as *Ca. aciculata*, *Ca. honghensis*, *Ca. lantauensis*, *Ca. pseudoturangicola*, *Ca. pseudoyunnanensis*, and *Ca. yunnanensis* spp. nov. Overall, the results reflect a high diversity of *Calonectria* species in China.

**Article info:** Submitted: 7 June 2017; Accepted: 1 October 2017; Published: 17 October 2017.

## INTRODUCTION

The genus *Calonectria* (*Hypocreales*, *Nectriaceae*) includes numerous important pathogens that cause significant damage to a large number of herbaceous and woody plants worldwide (Crous 2002, Lombard et al. 2010a). Approximately 335 plant species residing in about 100 plant families are hosts to *Calonectria* species, including important plantation tree crops such as species of *Eucalyptus*, *Pinus*, and *Acacia* (Crous 2002, Lombard et al. 2010a). To date, at least 149 species of *Calonectria* have been described and verified based on comparisons of DNA sequence data (Lombard et al. 2016, Marin-Felix et al. 2017). *Calonectria* species are soil-borne fungi (Thies & Patton 1970, Hwang & Ko 1976, Gilligan 1983, Crous 2002) and disease symptoms resulting from infection include cutting rot, damping-off, leaf blight, red crown rot, root rot, seedling rot, shoot blight and stem canker (Crous et al. 1991, Brown & Ferreira 2000, Crous 2002, Lombard et al. 2010a, 2015).

In China, plantation forestry utilizing rapidly-growing *Eucalyptus* species has expanded during the course of the

past two decades, to meet an increasing need for wood products. Approximately 4.5 M ha of *Eucalyptus* plantations have been established in South China (Chen & Chen 2013) and these are threatened by disease and insect pest problems (Zhou et al. 2008). Recent surveys of *Eucalyptus* plantations in South China have recorded several important emerging diseases, which include stem diseases caused by *Teratosphaeria zuluensis* (Cortinas et al. 2006, Chen et al. 2011a), species of *Botryosphaeriaceae* (Chen et al. 2011d) and *Cryphonectriaceae* (Chen et al. 2010, 2011b), and also *Ceratocystis* species (Chen et al. 2013). Leaf and shoot diseases caused by species of *Mycosphaerellaceae* and *Teratosphaeriaceae* (Burgess et al. 2006, 2007), *Quambalaria* species (Zhou et al. 2007), and *Calonectria* species (Crous et al. 2004, Lombard et al. 2010d, 2015, Chen et al. 2011c) have become widespread. Of these, *Calonectria* associated diseases are considered amongst the most threatening.

Pathogenic *Calonectria* species can cause significant losses to the *Eucalyptus* industry in China. The most important factor contributing to *Calonectria* infection and disease development is high humidity and free moisture (Crous 2002,

## Key words:

*Cylindrocladium*  
forest pathogens  
*Nectriaceae*  
phylogeny  
soil  
systematics

© 2017 International Mycological Association

You are free to share - to copy, distribute and transmit the work, under the following conditions:

**Attribution:** You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

**Non-commercial:** You may not use this work for commercial purposes.

**No derivative works:** You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.

Rodas et al. 2005). Common conditions in many parts of China where *Eucalyptus* species are propagated, serious disease problems emerge.

Twenty-eight species of *Calonectria* have been identified in China (Crous et al. 2004, Lombard et al. 2010d, 2015, Chen et al. 2011c, Xu et al. 2012). With the exception of *Ca. nymphaeae* (Xu et al. 2012), all species have been isolated from leaves, seedlings, and soil collected in *Eucalyptus* plantations and nurseries in South China (Supplementary Table 1). Twelve of these species were isolated from symptomatic *Eucalyptus* tissues, 17 were reported from soil associated with *Eucalyptus* trees or in *Eucalyptus* nurseries, and *Ca. pentaseptata* and *Ca. terrestris* were isolated from symptomatic *Eucalyptus* tissues as well as soil (Crous et al. 2004, Lombard et al. 2010d, 2015, Chen et al. 2011c). Pathogenicity tests have shown that 15 *Calonectria* species, including four known only from soil, are pathogenic to two tested *E. urophylla* × *E. grandis* hybrid clones commonly planted in South China (Chen et al. 2011c, Li et al. 2014a, b). Nothing is known regarding the pathogenicity of the remaining 11 *Calonectria* species known only from soil (Supplementary Table 1).

Species of *Calonectria* are characterised by a sexual morph having yellow to dark red perithecia, scaly to warty ascocarpial walls and 4–8-spored clavate ascii. The asexual morphs produce branched conidiophores, cylindrical, septate conidia and stipe extensions with terminal vesicles of characteristic shape (Crous 2002, Lombard et al. 2010c, 2016). These asexual morphs provide the best diagnostic characters for identification, especially in conidial and vesicle morphology (Schoch et al. 2000, Crous 2002). Based on phylogenetic inference that matches with the distribution of vesicle shapes, species of *Calonectria* are divided into two main groups. These include the Prolate Group including species with clavate to pyriform to ellipsoidal vesicles and the Sphaero-Naviculate Group that accommodates species with sphaeropedunculate and naviculate vesicles (Lombard et al. 2010c). At present, 14 species of *Calonectria* found in China reside in the Prolate Group and these include four species complexes: the *Ca. candelabrum* complex (*Ca. pauciramosa*, *Ca. seminaria*, and *Ca. tetraramosa*), *Ca. colhounii* complex (*Ca. fujianensis*, *Ca. nymphaeae*, and *Ca. pseudocolhounii*), *Ca. cylindrospora* complex (*Ca. cerciana*, *Ca. folicola*, *Ca. papillata*, and *Ca. terrestris*), and *Ca. reteaudii* complex (*Ca. crouxiana*, *Ca. pentaseptata*, *Ca. microconidialis*, and *Ca. pseudoreteaudii*) (Supplementary Table 1). The remaining 14 known species in China reside in the Sphaero-Naviculate Group and they all cluster in the *Ca. kyotensis* complex (Crous et al. 2004, Lombard et al. 2010d, 2015, Chen et al. 2011c, Xu et al. 2012) (Supplementary Table 1).

Previous research has suggested a relatively high *Calonectria* species diversity in South China (Chen et al. 2011c, Lombard et al. 2015). This study was undertaken in order to provide a more comprehensive overview of *Calonectria* species associated with planted *Eucalyptus* in the provinces of South China.

## MATERIALS AND METHODS

### Isolates

Surveys for *Calonectria* species were conducted in *Eucalyptus* plantations and nurseries of the Fujian, GuangDong, GuangXi, and YunNan Provinces in South China (Table 1). Leaves on trees showing blight symptoms were collected in *Eucalyptus* plantations. In *Eucalyptus* nurseries, seedlings showing stem and leaf rot symptoms were selected. Soil in the *Eucalyptus* plantations, and soil samples or planting substrate in *Eucalyptus* nurseries, were also sampled. In addition, soil samples were collected in a naturally forested area on Lantau Island in Hong Kong (Table 1). At each sampling site, between five and 25 *Eucalyptus* trees or seedlings were sampled, and between 10 and 25 soil samples were collected between March 2014 and May 2015. The symptomatic tissues were incubated in moist chambers at room temperature for 1–7 d to induce *Calonectria* sporulation. Soil samples were baited with germinating *Medicago sativa* (alfalfa) seeds using the method described by Crous (2002).

Conidial masses were transferred directly from *Eucalyptus* or *M. sativa* infected tissues to 2 % (v/v) malt extract agar (MEA) under a AxioCam Stemi 2000C dissecting microscope (Carl Zeiss, Germany). After incubation at room temperature for 2–5 d, a single hyphal tip from each culture was transferred to MEA plates and incubated at room temperature for 1 wk to obtain pure cultures.

Cultures were deposited in the Culture Collection of the China Eucalypt Research Centre (CERC), Chinese Academy of Forestry (CAF), ZhanJiang, GuangDong Province, China, and in the culture collection (CMW) of the Forestry Agricultural and Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa. Representative isolates including the ex-type cultures were deposited in the culture collection (CBS) of the Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands. Dried specimens of sporulating cultures were deposited in the National Collection of Fungi (PREM), Pretoria, South Africa.

### DNA extraction, PCR and sequence reactions

Isolates from sampled trees, seedlings and soil representing all sampling sites were used for total genomic DNA extraction and sequence comparisons. The DNA was extracted from 5–7-d-old cultures, using the CTAB method “5” as described by Van Burik et al. (1998). DNA concentrations were determined using a NanoDrop ND-2000 Spectrometer (Thermo Fisher Scientific, Waltham, MA). Four gene regions including the partial β-tubulin (*tub2*), calmodulin (*cmdA*), histone H3 (*his3*) and translation elongation factor 1-alpha (*tef1*) were amplified using the primers and protocols described by Lombard et al. (2010c). The TopTaqTM Master Mix Kit (Qiagen, Hilden) was used to amplify these gene regions. All PCR products were sequenced in both directions, using the same primers used for PCR amplification by the Beijing Genomics Institution, Guangzhou, China. All sequences obtained in this study were edited using Geneious v. 7.0 (Kearse et al. 2012) and were deposited in GenBank (Table 1).

### Phylogenetic analyses

All sequences representing the different *Calonectria* species in this study were used together with published

Table 1. Species of *Calonectria* collected in this study.

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>		
						<i>tef1</i>	<i>his3</i>	<i>cmaA</i>
<i>Ca. aciculata</i>	CMW 47645 <sup>5-6</sup> ; CERC 5342; CBS 142883	AAAA	Eucalyptus urophylla × <i>E. grandis</i> leaf in plantation	WeiYuan, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442644	MF442759	MF442874
<i>Ca. arbusta</i>	CMW 47502; CERC 9516	AAA <sup>-9</sup>	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442645	MF442760	MF442875
	CMW 47503; CERC 9520	AAA-	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442646	MF442761	MF442876
	CMW 47504; CERC 9522	AAA-	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442647	MF442762	MF442877
	CMW 47505; CERC 9523	AAA-	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442648	MF442763	MF442878
	CMW 47506; CERC 9525	AAA-	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442649	MF442764	MF442879
	CMW 47507; CERC 9526	AAA-	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442650	MF442765	MF442880
	CMW 47508 <sup>5</sup> ; CERC 9527	AAA-	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442651	MF442766	MF442881
	CMW 47509; CERC 9528	AAA-	Soil in <i>Eucalyptus</i> plantation	XiaoPingYang, XingBin, LaiBin, GuangXi, China	S.B. Liang	MF442652	MF442767	MF442882
	CMW 47637 <sup>5</sup> ; CERC 5320	AAA-	Soil in <i>Eucalyptus</i> plantation	XiDi, LongXu, WuZhou, GuangXi, China	S.F. Chen, J.Q. Li & W. Lu	MF442653	MF442768	MF442883
	CMW 47638; CERC 5322	AAA-	Soil in <i>Eucalyptus</i> plantation	XiDi, LongXu, WuZhou, GuangXi, China	S.F. Chen, J.Q. Li & W. Lu	MF442654	MF442769	MF442884
	CMW 47639; CERC 5324	AAA-	Soil in <i>Eucalyptus</i> plantation	XiDi, LongXu, WuZhou, GuangXi, China	S.F. Chen, J.Q. Li & W. Lu	MF442655	MF442770	MF442885
<i>Ca. asiatica</i>	CMW 47641 <sup>5</sup> ; CERC 5333	AAA-	Soil in <i>Eucalyptus</i> plantation	ZhengXing, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442656	MF442771	MF442886
	CMW 47654 <sup>5</sup> ; CERC 5373	ABB-	Soil in <i>Eucalyptus</i> plantation	WeiYuan, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442657	MF442772	MF442887
	CMW 47256 <sup>5</sup> ; CERC 3339	AAAA	Soil	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442658	MF442773	MF442888
	CMW 47258 <sup>5</sup> ; CERC 3349	ABAA	Soil	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442659	MF442774	MF442889
	CMW 47259 <sup>5</sup> ; CERC 3350	ABAA	Soil	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442660	MF442775	MF442890
	CMW 47260; CERC 3351	ABAA	Soil	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442661	MF442776	MF442891

Table 1. (Continued).

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>		
						<i>tef1</i>	<i>his3</i>	<i>cmda</i>
<i>Ca. eucalyptii</i>	CMW 47660 <sup>b</sup> ; CERC 5401	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	WeiYuan, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442662	MF442777	MF442892
<i>Ca. honghensis</i>	CMW 47667 <sup>b</sup> ; CERC 5568	AAAAA	Soil in <i>Eucalyptus</i> plantation	XinXian, PingBian, HongHe, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442663	MF442778	MF442893
	CMW 47668 <sup>b-7</sup> ; CBS 142884	AAAAA	Soil in <i>Eucalyptus</i> plantation	XinXian, PingBian, HongHe, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442664	MF442779	MF442894
	CMW 47669 <sup>b-8</sup> ; CBS 142885	AAAAA	Soil in <i>Eucalyptus</i> plantation	XinXian, PingBian, HongHe, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442665	MF442780	MF442895
	CMW 47670 <sup>b-7</sup> ; CBS 142886	AAAAA	Soil in <i>Eucalyptus</i> plantation	XinXian, PingBian, HongHe, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442666	MF442781	MF442896
	CMW 47671 <sup>b</sup> ; CERC 5574	AAAAA	Soil in <i>Eucalyptus</i> plantation	XinXian, PingBian, HongHe, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442667	MF442782	MF442897
<i>Ca. hongkongensis</i>	CMW 47257 <sup>b</sup> ; CERC 3341	AAAAA	Soil	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442668	MF442783	MF442898
	CMW 47271; CERC 3570	AAAAA	Soil in <i>Eucalyptus</i> plantation	ChangLe, HePu, BeiHai, GuangXi, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442669	MF442784	MF442899
	CMW 47274 <sup>b</sup> ; CERC 3573	AAAAA	Soil in <i>Eucalyptus</i> plantation	ChangLe, HePu, BeiHai, GuangXi, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442670	MF442785	MF442900
	CMW 47495; CERC 7125	AAAAA	Soil	FAFU, CangShan, FuZhou, Fujian, China	S.F. Chen	MF442671	MF442786	MF442901
	CMW 47499; CERC 7132	AAAAA	Soil	FAFU, CangShan, FuZhou, Fujian, China	S.F. Chen	MF442672	MF442787	MF442902
	CMW 47500; CERC 7133	AAAAA	Soil	FAFU, CangShan, FuZhou, Fujian, China	S.F. Chen	MF442673	MF442788	MF442903
	CMW 47501; CERC 7137	AAAAA	Soil	FAFU, CangShan, FuZhou, Fujian, China	S.F. Chen	MF442674	MF442789	MF442904
	CMW 47619; CERC 3288	AAA—	Soil	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442675	MF442790	MF442905
<i>Ca. lantauensis</i>	CMW 47251 <sup>b-7</sup> ; CBS 142887	AAA—	Soil	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442676	MF442791	MF442906
	CMW 47252 <sup>b-8</sup> ; CBS 142888	AAAAA	Medium of <i>E. urophylla</i> × <i>E. grandis</i> seedling in nursery	Lantau, Lidao, Hong Kong, China	M.J. Wingfield & S.F. Chen	MF442677	MF442792	MF442907
<i>Ca. mossambicensis</i>	CMW 47465 <sup>b</sup> ; CERC 6979	AAAAA	Medium of <i>E. urophylla</i> × <i>E. grandis</i> seedling in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442678	MF442793	MF442908
	CMW 47466; CERC 6990	AAAAA	Medium of <i>E. urophylla</i> × <i>E. grandis</i> seedling in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442679	MF442794	MF442909

Table 1. (Continued).

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>			
						tef1	his3	cmda	
								tub2	
CMW 47467; CERC 6996	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442680	MF442795	MF442910	MF443010	
CMW 47469; CERC 7004	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442681	MF442796	MF442911	MF443011	
CMW 47472; CERC 7022	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442682	MF442797	MF442912	MF443012	
CMW 47476; CERC 7038	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442683	MF442798	MF442913	MF443013	
CMW 47478; CERC 7048	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442684	MF442799	MF442914	MF443014	
CMW 47479; CERC 7056	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442685	MF442800	MF442915	MF443015	
CMW 47481; CERC 7072	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442686	MF442801	MF442916	MF443016	
CMW 47484 <sup>b</sup> ; CERC 7085	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442687	MF442802	MF442917	MF443017	
Ca. <i>pentaseptata</i>	CMW 47261; CERC 3529	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442688	MF442803	MF442918	MF443018
	CMW 47262; CERC 3533	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442689	MF442804	MF442919	MF443019
	CMW 47263; CERC 3535	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442690	MF442805	MF442920	MF443020
	CMW 47264; CERC 3536	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442691	MF442806	MF442921	MF443021
	CMW 47265; CERC 3537	AAAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442692	MF442807	MF442922	MF443022

Table 1. (Continued).

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>
					terf1 his3 cmndA	tub2
CMW 47266; CERC 3542	CERC	AAAA	Soil in <i>Eucalyptus</i> nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442693 MF442808 MF442923 MF443023
CMW 47267; CERC 3552	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442694 MF442809 MF442924 MF443024
CMW 47268; CERC 3559	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling medium in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442695 MF442810 MF442925 MF443025
CMW 47269; CERC 3560	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling medium in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442696 MF442811 MF442926 MF443026
CMW 47270 <sup>b</sup> ; CERC 3565	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & J.Q. Li	MF442697 MF442812 MF442927 MF443027
CMW 47277 <sup>b</sup> ; CERC 3652	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442698 MF442813 MF442928 MF443028
CMW 47278; CERC 3655	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442699 MF442814 MF442929 MF443029
CMW 47279; CERC 3658	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442700 MF442815 MF442930 MF443030
CMW 47280; CERC 3660	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442701 MF442816 MF442931 MF443031
CMW 47281; CERC 3664	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442702 MF442817 MF442932 MF443032
CMW 47282; CERC 3672	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442703 MF442818 MF442933 MF443033
CMW 47283; CERC 3680	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442704 MF442819 MF442934 MF443034
CMW 47284; CERC 3708	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442705 MF442820 MF442935 MF443035

Table 1. (Continued).

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>		
						tef1	his3	cmdA
CMW 47285; CERC 3720	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442706	MF442821	MF442936	MF443036
CMW 47463; CERC 6963	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling leaf in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & G.Q. Li	MF442707	MF442822	MF442937	MF443037
CMW 47464; CERC 6973	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling leaf in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen & G.Q. Li	MF442708	MF442823	MF442938	MF443038
CMW 47468; CERC 6999	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442709	MF442824	MF442939	MF443039
CMW 47470; CERC 7012	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442710	MF442825	MF442940	MF443040
CMW 47471; CERC 7018	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442711	MF442826	MF442941	MF443041
CMW 47473; CERC 7024	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442712	MF442827	MF442942	MF443042
CMW 47474; CERC 7030	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442713	MF442828	MF442943	MF443043
CMW 47475; CERC 7036	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442714	MF442829	MF442944	MF443044
CMW 47477; CERC 7047	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442715	MF442830	MF442945	MF443045
CMW 47480; CERC 7060	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442716	MF442831	MF442946	MF443046
CMW 47482; CERC 7074	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442717	MF442832	MF442947	MF443047
CMW 47483; CERC 7081	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442718	MF442833	MF442948	MF443048

Table 1. (Continued).

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>			
						ter1	hs3	cmdA	tub2
CMW 47485; CERC 7087	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442719	MF442834	MF442949	MF443049
CMW 47486; CERC 7095	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442720	MF442835	MF442950	MF443050
CMW 47487; CERC 7104	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> seedling stem in nursery	LingBei, SuiXi, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442721	MF442836	MF442951	MF443051
CMW 47510 <sup>5</sup> ; CERC 9529	CERC	AAAA	<i>Eucalyptus</i> clone seedling stem in nursery	CERC, XiaShan, ZhanJiang, GuangDong, China	J.Q. Li & S.F. Chen	MF442722	MF442837	MF442952	MF443052
CMW 47511; CERC 9533	CERC	AAAA	<i>Eucalyptus</i> clone seedling stem in nursery	CERC, XiaShan, ZhanJiang, GuangDong, China	J.Q. Li & S.F. Chen	MF442723	MF442838	MF442953	MF443053
CMW 47512; CERC 9541	CERC	AAAA	<i>Eucalyptus</i> clone seedling stem in nursery	CERC, XiaShan, ZhanJiang, GuangDong, China	J.Q. Li & S.F. Chen	MF442724	MF442839	MF442954	MF443054
CMW 47513; CERC 9556	CERC	AAAA	<i>Eucalyptus</i> clone seedling leaf in nursery	CERC, XiaShan, ZhanJiang, GuangDong, China	J.Q. Li & S.F. Chen	MF442725	MF442840	MF442955	MF443055
CMW 47514; CERC 9565	CERC	AAAA	<i>Eucalyptus</i> clone seedling stem in nursery	CERC, XiaShan, ZhanJiang, GuangDong, China	J.Q. Li & S.F. Chen	MF442726	MF442841	MF442956	MF443056
CMW 47515; CERC 9572	CERC	AAAA	<i>Eucalyptus</i> clone seedling stem in nursery	CERC, XiaShan, ZhanJiang, GuangDong, China	J.Q. Li & S.F. Chen	MF442727	MF442842	MF442957	MF443057
CMW 47620; CERC 3722	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442728	MF442843	MF442958	MF443058
CMW 47621; CERC 3730	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> stem in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442729	MF442844	MF442959	MF443059
CMW 47622; CERC 3736	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442730	MF442845	MF442960	MF443060
CMW 47623; CERC 3742	CERC	AAAA	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442731	MF442846	MF442961	MF443061

Table 1. (Continued).

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>			
						tef1	his3	cmaA	tub2
CMW 47624; CERC 3752	AAAA	E. urophylla × E. grandis leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442732	MF442847	MF442962	MF443062	
CMW 47625; CERC 3758	AAAA	E. urophylla × E. grandis leaf in plantation	HengShan, LianJiang, ZhanJiang, GuangDong, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442733	MF442848	MF442963	MF443063	
CMW 47626; CERC 4987	AAAA	E. urophylla × E. grandis leaf in plantation	JiuHe, Zijin, HeYuan, GuangDong, China	S.F. Chen & J.Q. Li	MF442734	MF442849	MF442964	MF443064	
CMW 47627; CERC 4989	AAAA	E. urophylla × E. grandis leaf in plantation	JiuHe, Zijin, HeYuan, GuangDong, China	S.F. Chen & J.Q. Li	MF442735	MF442850	MF442965	MF443065	
CMW 47628 <sup>b</sup> ; CERC 4992	AAAA	E. urophylla × E. grandis leaf in plantation	JiuHe, Zijin, HeYuan, GuangDong, China	S.F. Chen & J.Q. Li	MF442736	MF442851	MF442966	MF443066	
CMW 47629; CERC 4994	AAAA	E. urophylla × E. grandis leaf in plantation	JiuHe, Zijin, HeYuan, GuangDong, China	S.F. Chen & J.Q. Li	MF442737	MF442852	MF442967	MF443067	
CMW 47630; CERC 5005	AAAA	E. urophylla × E. grandis leaf in plantation	TongHe, PingNan, GuiGang, GuangXi, China	S.F. Chen & J.Q. Li	MF442738	MF442853	MF442968	MF443068	
CMW 47631 <sup>b</sup> ; CERC 5009	AAAA	E. urophylla × E. grandis leaf in plantation	TongHe, PingNan, GuiGang, GuangXi, China	S.F. Chen & J.Q. Li	MF442739	MF442854	MF442969	MF443069	
CMW 47632; CERC 5022	AAAA	E. urophylla × E. grandis leaf in plantation	TongHe, PingNan, GuiGang, GuangXi, China	S.F. Chen & J.Q. Li	MF442740	MF442855	MF442970	MF443070	
CMW 47633; CERC 5307	AAAA	E. urophylla × E. grandis leaf in plantation	ChengYue, SuiXi, ZhanJiang GuangDong, China	S.F. Chen & J.Q. Li	MF442741	MF442856	MF442971	MF443071	
CMW 47634; CERC 5310	AAAA	E. urophylla × E. grandis leaf in plantation	ChengYue, SuiXi, ZhanJiang GuangDong, China	S.F. Chen & J.Q. Li	MF442742	MF442857	MF442972	MF443072	
CMW 47635 <sup>b</sup> ; CERC 5313	AAAA	E. urophylla × E. grandis leaf in plantation	ChengYue, SuiXi, ZhanJiang GuangDong, China	S.F. Chen & J.Q. Li	MF442743	MF442858	MF442973	MF443073	
CMW 47636; CERC 5317	AAAA	E. urophylla × E. grandis leaf in plantation	ChengYue, SuiXi, ZhanJiang GuangDong, China	S.F. Chen & J.Q. Li	MF442744	MF442859	MF442974	MF443074	
<i>Ca. pseudoturangicola</i>	CMW 47247 <sup>b</sup> ; CERC 3250	AAAA	Soil	Lantau, Lidao, Hong Kong, China	MF442745	MF442860	MF442975	MF443075	
	CMW 47248 <sup>b</sup> ; CERC 3251	AAAA	Soil	Lantau, Lidao, Hong Kong, China	MF442746	MF442861	MF442976	MF443076	

Table 1. (Continued).

Species <sup>1</sup>	Isolate No. <sup>2</sup>	Haplotype <sup>3</sup>	Substrate	Sampling site	Collector	GenBank accession No. <sup>4</sup>
					<i>ter1</i>	<i>his3</i>
					<i>cmdA</i>	<i>tub2</i>
<i>CMW 47488<sup>5</sup>; CERC 7111</i>		AAAA	Soil in <i>Eucalyptus</i> plantation	BaiSha, MinHou, FuZhou, Fujian, China	S.F. Chen	MF442747 MF442862 MF442977 MF443077
<i>CMW 47489<sup>5-7</sup>; CERC 7115; CBS 142889</i>		AAAA	Soil in <i>Eucalyptus</i> plantation	BaiSha, MinHou, FuZhou, Fujian, China	S.F. Chen	MF442748 MF442863 MF442978 MF443078
<i>CMW 47490<sup>5</sup>; CERC 7116</i>		AAAA	Soil in <i>Eucalyptus</i> plantation	BaiSha, MinHou, FuZhou, Fujian, China	S.F. Chen	MF442749 MF442864 MF442979 MF443079
<b>CMW 47496<sup>5-8</sup>; CERC 7126; CBS 142890</b>		AAAA	Soil	FAFU, CangShan, FuZhou, Fujian, China	S.F. Chen	MF442750 MF442865 MF442980 MF443080
<i>CMW 47497<sup>5-7</sup>; CERC 7127; CBS 142891</i>		AAAB	Soil	FAFU, CangShan, FuZhou, Fujian, China	S.F. Chen	MF442751 MF442866 MF442981 MF443081
<i>CMW 47498<sup>5</sup>; CERC 7131</i>		AAAB	Soil	FAFU, CangShan, FuZhou, Fujian, China	S.F. Chen	MF442752 MF442867 MF442982 MF443082
<b>Ca. pseudoyunnanensis</b>	<b>CMW 47655<sup>5-8</sup>; CERC 5376; CBS 142892</b>	AAAA	Soil in <i>Eucalyptus</i> plantation	WeiYuan, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442753 MF442868 MF442983 MF443083
	<i>CMW 47656<sup>5-7</sup>; CERC 5377; CBS 142893</i>	AAAA	Soil in <i>Eucalyptus</i> plantation	WeiYuan, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442754 MF442869 MF442984 MF443084
	<i>CMW 47657<sup>5-7</sup>; CERC 5378; CBS 142894</i>	AAAA	Soil in <i>Eucalyptus</i> plantation	WeiYuan, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442755 MF442870 MF442985 MF443085
	<i>CMW 47642<sup>5-7</sup>; CERC 5337; CBS 142895</i>	AAAA	Soil in <i>Eucalyptus</i> plantation	ZhengXing, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442756 MF442871 MF442986 MF443086
	<i>CMW 47643<sup>5-7</sup>; CERC 5338; CBS 142896</i>	AAAA	Soil in <i>Eucalyptus</i> plantation	ZhengXing, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442757 MF442872 MF442987 MF443087
<b>Ca. yunnanensis</b>	<b>CMW 47644<sup>5-8</sup>; CERC 5339; CBS 142897</b>	AAAA	Soil in <i>Eucalyptus</i> plantation	ZhengXing, JingGu, PuEr, YunNan, China	S.F. Chen, J.Q. Li & G.Q. Li	MF442758 MF442873 MF442988 MF443088

<sup>1</sup> New species described in this study are indicated in bold.<sup>2</sup> CERC: China Eucalypt Research Centre, Zhanjiang, GuangDong Province, China; CMW: culture collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa. CBS: culture collection of Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands.<sup>3</sup> Haplotype within each identified species, determined by sequences of *ter1*, *his3*, *cmdA* and *tub2* regions.<sup>4</sup> *ter1* = translation elongation factor 1-alpha; *his3* = histone H3; *cmdA* = calmodulin; *tub2* = β-tubulin.<sup>5</sup> Isolates used in phylogenetic analyses.<sup>6</sup> Isolates used in morphological studies.<sup>7</sup> Isolates used in growth studies.<sup>8</sup> Isolates that represent ex-type cultures are indicated in bold.<sup>9</sup> “—” represents sequences that are not available.

sequences from ex-type strains of *Calonectria* downloaded from NCBI (<http://www.ncbi.nlm.nih.gov>) and subjected to phylogenetic analyses. Sequences generated in this study and those from NCBI were aligned using the online version of MAFFT v. 7 (<http://mafft.cbrc.jp/alignment/server>; Katho & Standley 2013) with the interactive refinement method (FFT-NS-i) setting. The aligned sequences were edited manually in MEGA v. 6 (Tamura *et al.* 2013) where necessary and deposited in TreeBASE (<http://treebase.org>). Single nucleotide polymorphisms (SNPs) were determined for each gene region between novel species identified in this study and their phylogenetically closest related species.

Based on sequences for *cmdA*, *his3*, *tef1* and *tub2* gene regions, the haplotypes of obtained *Calonectria* isolates were determined. Isolates representing different haplotypes and representing all the sampling sites were selected for the phylogenetic analyses. For the new species identified here, all the isolates were included in the analyses. The datasets were separated into two groups based on morphological characteristics representing the Prolate Group and the Sphaero-Naviculate Group, as defined by Lombard *et al.* (2010c). Phylogenetic analyses were conducted separately on the datasets for each of the four gene regions and combined data for three or four gene regions for the two groups, depending on the availability of *tub2* sequences for the *Calonectria* species selected for the phylogenetic analyses. A partition homogeneity test (PHT) was used to test whether conflict existed between the different datasets, the sequence data for coding gene regions were combined if no significant conflict (Cunningham 1997, Dettman *et al.* 2003). Data were analysed using Maximum Parsimony (MP) with PAUP\* v. 4.0b10 (Swofford 2003) and Maximum likelihood (ML) with PhyML v. 3.0 (Guindon & Gascuel 2003).

For MP analyses, gaps were treated as a fifth character (Ogden & Rosenberg 2007) and the characters were unordered and of equal weight with 1000 random addition replicates. The most parsimonious trees were generated using the heuristic search option with random stepwise addition of 1000 replicates and tree bisection and reconstruction (TBR) branch swapping. Zero-length branches were collapsed. Statistical support for internal nodes in trees was set with 1000 bootstrap replicates. Statistics estimated for parsimony included tree length (TL), retention index (RI), consistency index (CI), rescaled consistency indexes (RC) and homoplasy index (HI) (Hillis & Bull 1993).

For ML analyses, the appropriate models were obtained with jModeltest v. 2.1.5 (Posada 2008). The maximum number of retained trees was set to 1000 and the confidence levels for node support were determined using non-parametric bootstrapping with 1000 replicates. *Calonectria hongkongensis* (CBS 114828 and CBS 114711) and *Ca. pauciramosa* (CMW 5683 and CMW 30823) were used as the outgroup taxa for the Prolate Group and Sphaero-Naviculate Group, respectively. For all the analyses, the phylogenetic trees were viewed using MEGA v. 6 (Tamura *et al.* 2013).

## Morphology

Isolates were examined to define the characteristics of the asexual sporing structures. Single hyphal tip isolates were transferred to synthetic nutrient-poor agar (SNA; Nirenberg

1981, Lombard *et al.* 2010b, c) and incubated at room temperature for 7–15 d. The structures were examined and recorded using a Zeiss Axio Imager A1 microscope and a Zeiss AxioCam MRc digital camera with Zeiss Axio Vision Rel. 4.8 software (Carl Zeiss, Munchen). Morphological characteristics were studied by mounting the structures in a drop of 85 % lactic acid on glass slides. For the known *Calonectria* species, the structures were compared with those published of the type specimens. For those species shown to represent novel phylogenetic species, for ascospores, asci, conidia, and vesicles, 50 measurements were made for the isolates selected to represent the holotype specimen. In addition, 30 measurements were made for paratype specimens. Minimum, maximum and average (mean) values were calculated and are presented as (minimum–) (average – standard deviation)–(average + standard deviation) (–maximum). For all other taxonomic informative structures, only the extremes are given.

The optimal growth conditions for cultures representing novel species were determined on MEA in the dark at temperatures ranging from 5–35 °C with 5 °C intervals. Four replicates were used for each isolate at each temperature. Two diameter measurements, perpendicular to each other, were measured daily for 7 d. Colony morphology and colour were determined on MEA after growth at 25 °C in the dark for 7 d using the colour charts of Rayner (1970). All descriptions were deposited in MycoBank ([www.mycobank.org](http://www.mycobank.org); Crous *et al.* 2004).

## Sexual compatibility

Isolates of each novel *Calonectria* species identified based on multi-gene phylogenetic analyses were crossed with each other in all possible combinations. Crosses were made on minimal salt agar (MSA) on which sterile toothpicks had been placed on the surface of the media (Geurber & Correll 2001, Lombard *et al.* 2010b, c) and incubated at room temperature. Isolates crossed with themselves served as controls, and it was thus possible to distinguish between those species with heterothallic or homothallic mating systems. Crosses were regarded as successful when isolate combinations produced perithecia extruding viable ascospores.

After 4–6 wk of incubation, the perithecia obtained from the sexual compatibility tests were mounted in Leica Biosystems Tissue Freezing Medium (Leica Biosystems Nussloch, Nussloch, Germany) and sectioned using a Microtome Cryostat Microm HM550 (Microm International, Thermo Fisher Scientific, Walldorf, Germany) at –20 °C to observe characteristics of the ascomata and ascostromatic tissues. The 12 µm sections were mounted in 85 % lactic acid and 3 % KOH, and all taxonomically informative structures were measured in the same manner as that for the asexual structures.

## RESULTS

### Isolates

A total of 115 isolates having morphological characteristics typical of *Calonectria* spp. were obtained. Of these, 64 isolates were from GuangDong Province, 16 from GuangXi

Province, 15 from YunNan Province, 10 from Fujian Province, and 10 isolates were from soil in a natural forested area in the Hong Kong Region. All of these isolates were either from soil samples (mostly from beneath *Eucalyptus* trees), from infected leaves on *Eucalyptus* trees or from *Eucalyptus* plants in nurseries (Table 1).

## Phylogenetic analyses

All 115 isolates obtained in this study were sequenced (Table 1). Thus, approximately 475 bp were generated for the *cmdA* gene region, 435 bp for the *his3* gene region, 500 bp for the *tef1* gene region and 565 bp for the *tub2* gene region. The 115 isolates represent 16 haplotypes determined by sequences for the four gene regions (Table 1). In total, 40 isolates collected in this study which represent all the 16 haplotypes were selected for phylogenetic analyses. Based on the comparisons for four gene region sequences generated in this study and published sequences from ex-type strains of *Calonectria* downloaded from NCBI, sequences for 65 ex-type and other strains representing 34 species closely related to species emerging from this study were used for analyses (Supplementary Table 2). For the 40 isolates selected for phylogenetic analyses, 15 resided in the Prolate Group and 25 isolates formed part of the Sphaero-Naviculate Group. For the MP and ML trees based on the single and combined sequence datasets (TreeBASE no 21167) in Prolate Group or Sphaero-Naviculate Group, although the relative positions of individual *Calonectria* species differed slightly, while the overall topologies were similar.

*Species residing in the Prolate Group:* The partition homogeneity tests (PHT) for combinations of the *tef1*, *his3*, *cmdA* and *tub2* gene regions yielded a P-value of 0.001, and consequently, the sequence data for coding gene regions were combined (Cunningham 1997, Dettman *et al.* 2003). The combined dataset included 51 taxa and consisted of 1993 characters, including alignment gaps, of which 1414 were parsimony-uninformative and 579 were parsimony-informative. Statistical values for the trees for the MP analyses and parameters for the best-fit substitution models of ML are provided in Supplementary Table 3. The ML tree of combined sequence dataset is presented in Fig. 1.

In total, the 15 isolates collected in this study residing in the Prolate Group clustered in three phylogenetic groups (Group A, Group B and Group C), which belong to the *Ca. colhounii*, *Ca. reteaudii* and *Ca. candelabrum* complexes, respectively (Fig. 1). In Group A, five isolates (CMW 47667, CMW 47668, CMW 47669, CMW 47670 and CMW 47671) grouped in a novel monophyletic cluster (ML/MP: 89 % / 92 %) with a single isolate, CMW 47645, forming a novel distinct basal lineage, both of two novel lineages were closely related to, but separate from *Ca. monticola* and *Ca. colhounii* (Fig. 1). The total number of the fixed unique differences (SNPs) between the four clades for all four gene regions combined varied between 12–26 (Supplementary Table 4). One isolate (CMW 47660) was identified as *Ca. eucalypti* (Fig. 1).


In Group B, six isolates (CMW 47270, CMW 47277, CMW 47510, CMW 47628, CMW 47631 and CMW 47635) resided in the same clade as *Ca. pentaseptata*. Two isolates (CMW 47465 and CMW 47484) clustered within the clade representing *Ca. mossambicensis* in Group C (Fig. 1).

*Species in the Sphaero-Naviculate Group:* For this Group, sequences for the *tub2* gene region were not available for some taxa due to multiple sequence copies occur in single *Calonectria* isolates. The PHT comparing the *tef1*, *his3* and *cmdA* gene regions gave a P = 0.077 value. This showed that there was no significant conflict between the three gene regions and the sequence data for three gene regions were combined (Cunningham 1997, Dettman *et al.* 2003). The combined sequence dataset included 58 taxa and consisted of 1415 characters, including alignment gaps. Of these, 1015 were parsimony-uninformative and 400 were parsimony-informative. Statistical values for the MP trees and parameters for the best-fit substitution models of ML are provided in Supplementary Table 3. The ML tree is presented in Fig. 2.

The 25 isolates placed in the Sphaero-Naviculate Group of *Calonectria* collected in this study clustered into three phylogenetic groups (Groups D–F), which all belong to the *Ca. kyotensis* complex (Fig. 2). Group D included six isolates residing in two distinct sister clades; CMW 47642, CMW 47643 and CMW 47644 in one clade, and CMW 47655, CMW 47656 and CMW 47657 in another clade (ML/MP: 85 % / 80 %, ML/MP: 71 % / 73 %, respectively). Three and four SNPs could be identified in each of the two clades for *his3* and *tub2* gene sequences (Supplementary Table 5). These two clades were phylogenetically most closely related to *Ca. asiatica* and *Ca. colombiensis* (Fig. 2). The total number of SNP differences between isolates in these two clades, *Ca. asiatica* and *Ca. colombiensis*, for all four gene regions combined, varied between 7–28 (Supplementary Table 5). Two isolates (CMW 47251 and CMW 47252) formed a single independent clade that was distinct from any known *Calonectria* species and this was supported by high bootstrap values (ML/MP: 100 % / 100 %) (Fig. 2). The total number of SNP differences between this clade accommodating isolates CMW 47251 and CMW 47252, and other phylogenetically closely related *Calonectria* species (*Ca. curvispora*, *Ca. illicicola*, *Ca. pacifica* and *Ca. sumatrensis*) for three gene regions combined varied between 15–44 (Supplementary Table 6). Isolates CMW 47641 and CMW 47654 resided in the clade representing *Ca. asiatica*, however, with low bootstrap support (Fig. 2). In addition, isolates CMW 47508 and CMW 47637 did not reside in a distinct clade but were closely related to *Ca. arbusta* (Fig. 2).

In Group E, eight isolates (CMW 47247, CMW 47248, CMW 47488, CMW 47489, CMW 47490, CMW 47496, CMW 47497 and CMW 47498) formed a well-resolved clade (ML/MP: 81 % / 98 %), close to, but distinct from *Ca. turangicola* (Fig. 2). Several SNPs could be identified for this clade and *Ca. turangicola*, for three of the four gene regions analysed (Supplementary Table 7). The total number of SNP differences between this clade and the species most closely related to it for all four gene regions combined, varied between 6–34 (Supplementary Table 7). Two isolates, CMW 47257 and CMW 47274 clustered with *Ca. hongkongensis* (Fig. 2).

In Group F, three isolates (CMW 47256, CMW 47258 and CMW 47259), representing two haplotypes, grouped in a clade, although, lacking bootstrap support. These isolates were most closely related to *Ca. chinensis* (Fig. 2).




**Fig. 1.** Phylogenetic tree based on ML analysis of a combined DNA dataset of *tef1*, *his3*, *cmdA* and *tub2* gene sequences for the species of *Calonectria* in the Prolate Group. Bootstrap value  $\geq 70\%$  for MP and ML analyses are presented at the branches. Bootstrap values lower than 70 % are marked with “\*”, and absent are marked with “-”. Isolates, representing ex-type material, are marked with “T”, isolates sequenced in this study are highlighted in blue and bold. The tree was rooted to *Ca. hongkongensis* (CBS 114828 and CBS 114711).

### Sexual compatibility

Sixteen isolates belonging to three of the novel taxa (*Ca. honghensis*, *Ca. pseudoturangicola*, and *Ca. yunnanensis*)

were able to produce sexual structures when crossed with themselves. These included isolates CMW 47247, CMW 47248, CMW 47488–47490, CMW 47496–47498,

## Sphaero-Naviculate Group



**Fig. 2.** Phylogenetic tree based on ML analysis of a combined DNA dataset of *tef1*, *his3* and *cmdA* gene sequences for the species of *Calonectria* in the Sphaero-Naviculate Group. Bootstrap value  $\geq 70\%$  for MP and ML analyses are presented at the branches. Bootstrap values lower than 70 % are marked with “\*”, and absent are marked with “-”. Isolates, representing ex-type material, are marked with “T”, isolates sequenced in this study are highlighted in blue and bold. The tree was rooted to *Ca. pauciramosa* (CMW 5683 and CMW 30823).

CMW 47642–47644 and CMW 47667–47671 that formed protoperithecia within 2–3 wk and perithecia within 4–6 wk. They were consequently recognised as homothallic. The remaining isolates identified as novel *Calonectria* species

failed to yield any perithecia in crosses, indicating that they were either self-sterile (heterothallic) or they lacked the ability to recombine to produce fertile progeny in culture.

## Morphology and taxonomy

Based on DNA sequence comparisons (Figs 1–2) and morphology, isolates collected in this study resided in either the Prolate or Sphaero-Naviculate Group of *Calonectria* species as defined by Lombard *et al.* (2010c). For the 40 isolates selected for phylogenetic analyses, 18 resolved as known species in six groups and respectively represented *Ca. eucalypti* (Group A; *Ca. colhounii* species complex), *Ca. pentaseptata* (Group B; *Ca. reteaudii* species complex), *Ca. mossambicensis* (Group C; *Ca. candelabrum* species complex), *Ca. asiatica* and *Ca. arbusta* (Group D), *Ca. hongkongensis* (Group E) and *Ca. chinensis* (Group F), the latter three groups all clustered in the *Ca. kyotensis* species complex (Figs 1–2). The former three species resided in the Prolate Group and the latter four known species all clustered in the Sphaero-Naviculate Group (Figs 1–2).

The remaining isolates grouped in six distinct clades (Figs 1–2) that represent novel taxa: *Calonectria aciculata* and *Ca. honghensis* spp. nov. in the Prolate Group; and *Ca. lantauensis*, *Ca. pseudoturangicola*, *Ca. pseudoyunnanensis*, and *Ca. yunnanensis* spp. nov. in Sphaero-Naviculate Group. The morphological characters of isolates identified as new species were compared with the species phylogenetically most closely related to them, and these characteristics are summarized in Table 2. Based on phylogenetic inference and morphological features, these isolates represent six previously undescribed species of *Calonectria* described below:

## TAXONOMY

### Species in the Prolate Group

#### *Calonectria aciculata* J.Q. Li, Q.L. Liu & S.F. Chen, sp. nov.

MycoBank MB821632  
(Fig. 3)

**Etymology:** After the acicular vesicles in this species.

**Diagnosis:** *Calonectria aciculata* can be distinguished from the phylogenetically closely related *Ca. colhounii*, *Ca. honghensis*, and *Ca. monticola* in the longer macroconidia.

**Type: China:** YunNan Province: PuEr Region, JingGu County, WeiYuan Town, on leaves of an *E. urophylla* × *E. grandis* hybrid clone, 16 Nov. 2014, S.F. Chen & J.Q. Li (PREM 61941 – holotype; CMW 47645 = CERC 5342 = CBS 142883 – ex-type cultures).

**Description:** Sexual morph unknown. Macroconidiophores consisting of a stipe, a suite of penicillate arranged fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 48–176 × 3–7 µm; stipe extensions septate, straight to flexuous 90–193 × 2.5–4 µm long, 2–4 µm wide at the apical septum, terminating in acicular to clavate vesicles, (2.0)–2.5–3.5(–5) µm diam. Conidiogenous apparatus 19–110 µm long, 27–145 µm wide; primary branches aseptate to 1-septate, 13–38 ×

3.5–6 µm; secondary branches aseptate, 11–24 × 3.5–5.5 µm; tertiary branches aseptate, 9–14 × 3.5–4.5 µm, each terminal branch producing 2–4 phialides; phialides doliform to reniform, hyaline, aseptate, 6–14 × 2.5–5 µm, apex with minute periclinal thickening and inconspicuous collarette. Macroconidia cylindrical, rounded at both ends, straight, (53)–62–76(–86) × (4.5)–5–6(–7) µm (av. = 69 × 5.5 µm), 3-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. Mega- and microconidia not observed.

**Culture characteristics:** Colonies forming abundant white aerial mycelium on MEA at 25 °C after 7 d; moderate sporulation, feathery, irregular margins, reverse pale ochraceous-salmon (13'f) to sanford's brown (11k). Chlamydospores common throughout the medium forming microsclerotia. **Growth characteristics,** optimal growth temperature 25 °C, no growth at 5 °C and 35 °C, after 7 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C and 30 °C reached 11.3 mm, 32.1 mm, 48.3 mm, 60.3 mm and 30.8 mm, respectively.

**Notes:** *Calonectria aciculata* differs from the phylogenetically closely related species *Ca. colhounii*, *Ca. honghensis*, and *Ca. monticola* with respect to the size of its macroconidia. The average sizes of the macroconidia of *Ca. aciculata* (av. = 69 × 5.5 µm) are longer than the average sizes of *Ca. colhounii* (av. = 65 × 5 µm), *Ca. honghensis* (av. = 54 × 5.5 µm) and *Ca. monticola* (av. = 49 × 5 µm) (Crous 2002, Crous *et al.* 2015b).

#### *Calonectria honghensis* J.Q. Li, Q.L. Liu & S.F. Chen, sp. nov.


MycoBank MB821633  
(Fig. 4)

**Etymology:** From the HongHe Region of China where the fungus was first collected.


**Diagnosis:** *Calonectria honghensis* differs from the phylogenetically closely related *Ca. aciculata*, *Ca. colhounii* and *Ca. monticola* in the dimensions of the macroconidia and ascospores.

**Type: China:** YunNan Province: HongHe Region, PingBian County, XinXian Town, from soil collected in a *Eucalyptus* plantation, 14 Nov. 2014, S.F. Chen & J.Q. Li (PREM 61943 – holotype; CMW 47669 = CERC 5572 = CBS 142885 – ex-type cultures).

**Description:** Perithecia solitary or in groups of up to four, yellow, becoming orange with age; in section apex and body yellow, base red-brown, subglobose to ovoid, 208–423 µm high, 233–406 µm diam, body turning dark yellow, and base dark red in 3 % KOH; perithecial walls rough consisting of two thick-walled layers: outside layer of *textura globulosa*, 10–57 µm wide, becoming more compressed towards inner layer of *textura angularis*, 10–23 µm wide, becoming thin-walled and hyaline towards the centre; outer cells 9–41 × 7–24 µm, inner cells 10–19



**Fig. 3.** *Calonectria aciculata*. **A–B.** Macroconidiophore. **C–E.** Acicular to clavate vesicles. **F–G.** Conidiogenous apparatus with conidiophore branches and doliiform to reniform phialides. **H–I.** Macroconidia. Bars: A–B = 50 µm; C–I = 10 µm.



**Fig. 4.** *Calonectria honghensis*. **A.** Perithecium. **B.** Vertical section through a perithecium. **C.** Cells around ostiolar region of perithecium. **D.** Section through lateral perithecial wall. **E–F.** Ascospores. **G.** Ascospores. **H–I.** Macroconidiophore. **J–K.** Clavate vesicles. **L–M.** Conidiogenous apparatus with conidiophore branches and doliform to reniform phialides. **N.** Macroconidia. Bars: A = 200 µm; B = 50 µm; C–F and H–I = 20 µm; G and L–N = 10 µm; J and K = 5 µm.

**Table 2.** Morphological comparisons of *Calonectria* species examined in this study and other phylogenetically closely related species.

Species	Ascospores (L × W) <sup>1,2</sup>	Ascospores average (L × W) <sup>1,2</sup>	Ascospores septation	Macroconidia (L × W) <sup>1,2</sup>	Macroconidia average (L × W) <sup>1,2</sup>	Macroconidia septation	Vesicle (Min.–Max.) <sup>3</sup>	Vesicle shape	Reference
<b><i>Ca. aciculata</i><sup>4</sup></b>	N/A <sup>5</sup>	N/A		(53–)62–76(–86) × (4.5–)5–6(–7) <sup>6</sup>	69 × 5.5	3	(2–)2.5–3.5(–5)	acicular to clavate	This study
<b><i>Ca. honghensis</i></b>	(35–)43–55(–65) × (4.5–)5.5–6.5(–7.5)	49 × 6	3	(43–)49–59(–66) × (4.5–)5–5.5(–6)	54 × 5.5	3	(2.5–)3–4.5(– 5.5)	clavate	This study
<b><i>Ca. colhounii</i></b>	(30–)50–65(–75) × (4–)5–6(–8)	55 × 6	(1–)3	(45–)60–70(–80) × (4–)5(–6)	65 × 5	(1–)3	3–4	clavate	Crous (2002)
<b><i>Ca. monticola</i></b>	N/A	N/A		46–51(–56) × 4–6(–7)	49 × 5	3	4–6	broadly clavate	Crous et al. (2015b)
<b><i>Ca. lantauensis</i></b>	N/A	N/A		(49–)52–58(–62) × (4.5–)5–5.5(–6)	55 × 5	1	(7.5–)8.5– 12.5(–17.5)	sphaeropedunculate	This study
<b><i>Ca. curvispora</i></b>	N/A	N/A		(45–)55–65(–70) × (4–)5–6	60 × 5	1(–3)	(5–)8(–10)	sphaeropedunculate	Crous (2002)
<b><i>Ca. illiticola</i></b>	(30–)37–50(–65) × (4–)5–6.5(–7)	45 × 6	1(–3)	(45–)70–82(–90) × (4–)5–6.5(–7)	62 × 6	(1–)3	(6–)7–10(–12)	sphaeropedunculate	Crous (2002)
<b><i>Ca. pacifica</i></b>	N/A	N/A		(38–)45–65(–75) × 4–5	55 × 4.5	1	7–15	sphaeropedunculate	Crous (2002)
<b><i>Ca. sumatrensis</i></b>	N/A	N/A		(45–)55–65(–70) × (4.5)(–6)	58 × 5	1	8–13	sphaeropedunculate	Crous et al. (2004)
<b><i>Ca. pseudoturangicola</i></b>	(24–)27–35(–43) × (4.5–)5.5–7.5(–9.5)	31 × 6.5	1(–3)	(33–)36–44(–50) × (2.5–)3.5–4	40 × 3.5	1	(4.5–)5–8.5(– 12)	sphaeropedunculate	This study
<b><i>Ca. hongkongensis</i></b>	(25–)28–35(–40) × (4–)5–6(–7)	31 × 6	1	(38–)45–48(–53) × (4–)4.5(–5)	46.5 × 4	1	8–14	sphaeropedunculate	Crous et al. (2004)
<b><i>Ca. turangicola</i></b>	N/A	N/A		(40–)42–46(–47) × 3–5	44 × 4	1	8–12	sphaeropedunculate	Lombard et al. (2015)
<b><i>Ca. pseudoyunnanensis</i></b>	N/A	N/A		(40–)44–50(–55) × (4–)4.5–5.5(–6)	47.5 × 5	1	(2.5–)3.5–5	ellipsoidal, obpyriform to sphaeropedunculate	This study
<b><i>Ca. yunnanensis</i></b>	(28–)31–41(–55) × (5–)5.5–6.5(–8)	36 × 6	1(–3)	(36–)39–47(–52) × (4–)4.5–5(–5.5)	43 × 4.5	1	(2–)2.5–3.5(– 4.5)	sphaeropedunculate	This study
<b><i>Ca. asiatica</i></b>	(28–)30–38(–40) × (5–)6(–7)	33 × 6	1	(42–)48–55(–65) × (4–)5(–5.5)	53 × 5	1	12–17	sphaeropedunculate	Crous et al. (2004)
<b><i>Ca. colombiensis</i></b>	(28–)30–35(–40) × (4–)5(–6)	33 × 5	1	(33–)48–58(–60) × (4–)4.5(–5)	53 × 4.5	1(–3)	7–12	sphaeropedunculate	Crous et al. (2004)

<sup>1</sup>All measurements are in µm.<sup>2</sup>L × W = length × width.<sup>3</sup>Min.–Max. = minimum–maximum.<sup>4</sup>Species indicated in bold are described in this study.<sup>5</sup>N/A = not available.<sup>6</sup>Measurements are presented in the format [(minimum–) (average – standard deviation)–(average + standard deviation) (–maximum)].

$\times$  3–13  $\mu\text{m}$ , perithecial base to 190  $\mu\text{m}$  wide, consisting of dark red, angular cells merging with an erumpent stroma, cells of the outer wall layer continuing into the pseudoparenchymatous cells of the erumpent stroma. Ascii 4-spored, clavate, (75–)91–115(–153)  $\times$  (13–)14–24(–37)  $\mu\text{m}$  (av. = 103  $\times$  19  $\mu\text{m}$ ), tapering to a long thin stalk. Ascospores aggregate in the upper third of the ascci, hyaline, guttulate, fusoid with rounded ends, straight to slightly curved, 3-septate, not or slightly constricted at the septum, (35–)43–55(–65)  $\times$  (4.5–)5.5–6.5(–7.5)  $\mu\text{m}$  (av. = 49  $\times$  6  $\mu\text{m}$ ). Macroconidiophores consisting of a stipe, a suite of penicillate arranged fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 42–192  $\times$  4–10  $\mu\text{m}$ ; stipe extensions septate, straight to flexuous, 70–215  $\mu\text{m}$  long, 3–5  $\mu\text{m}$  wide at the apical septum, terminating in a clavate vesicle, (2.5–)3.0–4.5(–5.5)  $\mu\text{m}$  diam. Conidiogenous apparatus 33–114  $\mu\text{m}$  long, 21–75  $\mu\text{m}$  wide; primary branches aseptate to 1-septate, 14–57  $\times$  4–7.5  $\mu\text{m}$ ; secondary branches aseptate, 10–26  $\times$  4–5.5  $\mu\text{m}$ ; tertiary branches aseptate, 9–19  $\times$  3.5–6  $\mu\text{m}$ ; additional branches (–4), aseptate, 9.5–14.5  $\times$  3.5–5  $\mu\text{m}$ ; each terminal branch producing 2–4 phialides; phialides doliform to reniform, hyaline, aseptate, 7–12  $\times$  3–5  $\mu\text{m}$ , apex with minute periclinal thickening and inconspicuous collarette. Macroconidia cylindrical, rounded at both ends, straight, (43–)49–59(–66)  $\times$  (4.5–)5–5.5(–6)  $\mu\text{m}$  (av. = 54  $\times$  5.5  $\mu\text{m}$ ), 3-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. Mega- and microconidia not observed.

**Culture characteristics:** Colonies forming white to sienna aerial mycelium on MEA at 25 °C after 7 d, profuse sporulation, feathery, irregular margins, reverse capucine buff (13f) to umber (9). Chlamydospores common throughout the medium forming microsclerotia. **Growth characteristics,** optimal growth temperature 25 °C, no growth at 5 °C and 35 °C, after 7 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C and 30 °C reached 14.3 mm, 31.6 mm, 43.5 mm, 51.9 mm and 17.3 mm, respectively.

**Additional material examined:** **China:** YunNan Province: HongHe Region, PingBian County, XinXian Town: from soil collected in a *Eucalyptus* plantation, 14 Nov. 2014, S.F. Chen & J.Q. Li (PREM 61942, culture CMW 47668 = CERC 5571 = CBS 142884; PREM 61944, culture CMW 47670 = CERC 5573 = CBS 142886).

**Notes:** *Calonectria honghensis* is phylogenetically most closely related to *Ca. aciculata*, *Ca. colhounii*, and *Ca. monticola*. However, *Ca. honghensis* can be distinguished from these species by the dimensions of the macroconidia and ascospores. The average size of the macroconidia of *Ca. honghensis* (av. = 54  $\times$  5.5  $\mu\text{m}$ ) is shorter than that of *Ca. aciculata* (av. = 69  $\times$  5.5  $\mu\text{m}$ ) and *Ca. colhounii* (av. = 65  $\times$  5  $\mu\text{m}$ ), but longer than that of *Ca. monticola* (av. = 49  $\times$  5  $\mu\text{m}$ ) (Crous 2002, Crous *et al.* 2015b). The average size of the ascospores of *Ca. honghensis* (av. = 49  $\times$  6  $\mu\text{m}$ ) is shorter than for *Ca. colhounii* (av. = 55  $\times$  6  $\mu\text{m}$ ) (Crous 2002); sexual structures are not known for *Ca. aciculata* and *Ca. monticola* (Crous *et al.* 2015b).

## Species in the Sphaero-Naviculate Group

**Calonectria lantauensis** J.Q. Li, Q.L. Liu & S.F. Chen, sp. nov.  
MycoBank MB821634  
(Fig. 5)

**Etymology:** After Lantau Island in Hong Kong, China, where the fungus was first collected.

**Diagnosis:** *Calonectria lantauensis* can be distinguished from the phylogenetically closely related species *Ca. curvispora*, *Ca. illicicola*, *Ca. pacifica* and *Ca. sumatreensis* by the size of the macroconidia.


**Type:** **China:** Hong Kong Region: LiDiao Distict, Lantau Island, from soil collected in roadside near Hong Kong airport, 12 Mar. 2014, M.J. Wingfield & S.F. Chen (PREM 61946 – holotype; CMW 47252 = CERC 3302 = CBS 142888 – ex-type cultures).

**Description:** Sexual morph unknown. Macroconidiophores consisting of a stipe, a suite of penicillate arranged fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 44–216  $\times$  4.5–12.5  $\mu\text{m}$ ; stipe extension septate, straight to flexuous 51–271  $\mu\text{m}$  long, 2–5.5  $\mu\text{m}$  wide at the apical septum, terminating in a sphaeropedunculate vesicle, (7.5–)8.5–12.5(–17.5)  $\mu\text{m}$  diam; lateral stipe extensions absent. Conidiogenous apparatus 45–173  $\mu\text{m}$  long, 34–114  $\mu\text{m}$  wide; primary branches aseptate to 1-septate, 16–83  $\times$  4.5–12.5  $\mu\text{m}$ ; secondary branches aseptate, 10–19  $\times$  4.5–7.5  $\mu\text{m}$ ; tertiary branches aseptate, 7.5–13  $\times$  3.5–6  $\mu\text{m}$ ; each terminal branch producing 2–6 phialides; phialides doliform to reniform, hyaline, aseptate, 5.5–13  $\times$  3–8  $\mu\text{m}$ ; apex with minute periclinal thickening and inconspicuous collarette. Macroconidia cylindrical, rounded at both ends, straight, (49–)52–58(–62)  $\times$  (4.5–)5–5.5(–6)  $\mu\text{m}$ , (av. = 55  $\times$  5  $\mu\text{m}$ ), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. Mega- and microconidia not observed.

**Culture characteristics:** Colonies forming abundant white to buff aerial mycelium on MEA at 25 °C after 7 d, moderate sporulation, feathery, irregular margins, reverse sienna (8) to umber (9). **Growth characteristic,** optimal growth temperature 25 °C, no growth at 5 °C and 35 °C, after 7 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C and 30 °C reached 7.2 mm, 28.8 mm, 54.2 mm, 78.0 mm and 71.6 mm, respectively.

**Additional material examined:** **China:** Hong Kong Region: LiDiao Distict, Lantau Island, from soil collected in roadside near Hong Kong airport, 12 Mar. 2014, M.J. Wingfield & S.F. Chen (PREM 61945, culture CMW 47251 = CERC 3301 = CBS 142887).

**Notes:** *Calonectria lantauensis* is closely related to *Ca. curvispora*, *Ca. illicicola*, *Ca. pacifica*, and *Ca. sumatreensis*. *Calonectria lantauensis* can be distinguished from *Ca. curvispora*, *Ca. illicicola* and *Ca. sumatreensis* by the average size of the macroconidia. The macroconidia of *Ca. lantauensis* (av. = 55  $\times$  5  $\mu\text{m}$ ) are shorter than those of



**Fig. 5.** *Calonectria lantauensis*. **A–B.** Macroconidiophore. **C–E.** Sphaeropedunculate vesicles. **F–G.** Conidiogenous apparatus with conidiophore branches and doliiiform to reniform phialides. **H–I.** Macroconidia. Bars: A = 50 µm; B and G = 20 µm; C–F and H–I = 10 µm.

*Ca. curvispora* (av. =  $60 \times 5$  µm), *Ca. illicicola* (av. =  $62 \times 6$  µm) and *Ca. sumatrensis* (av. =  $58 \times 5$  µm) (Crous 2002, Crous et al. 2004). No lateral stipe extensions were found in *Ca. lantauensis*, *Ca. curvispora* or *Ca. illicicola*, while these structures are commonly observed in *Ca. pacifica* but rarely observed in *Ca. sumatrensis* (Crous 2002, Crous et al. 2004).

**Calonectria pseudoturangicola** J.Q. Li, Q.L. Liu & S.F. Chen, sp. nov.

MycoBank MB821635  
(Fig. 6)

**Etymology:** From the close resemblance to *Calonectria turangicola*.

**Diagnosis:** *Calonectria pseudoturangicola* can be distinguished from the phylogenetically closely related species *Ca. hongkongensis* and *Ca. turangicola* in the shorter and narrower macroconidia.

**Type:** **China:** FuJian Province: FuZhou City, CangShan District, from soil collected in the campus of Fujian Agriculture and Forestry University (FAFU), 14 Dec. 2014, S.F. Chen (PREM 61948 – holotype; CMW 47496 = CERC 7126 = CBS 142890 – ex-type cultures).

**Description:** Perithecia solitary or in groups of up to five, orange, becoming orange-brown with age; in section, apex and body orange, base red-brown, subglobose to ovoid, 241–511 µm high, 242–456 µm diam, body turning red, and base dark red-brown in 3 % KOH; perithecial walls rough consisting of two thick-walled layers: outside layer of *textura globulosa*, 21–49 µm wide, becoming more compressed towards inner layer of *textura angularis*, 8–16 µm wide, becoming thin-walled and hyaline towards the centre; outer cells 17–60 × 10–33 µm, inner cells 7–44 × 2–15 µm; perithecial base to 191 µm wide, consisting of dark red, angular cells merging with an erumpent stroma, cells of the outer wall layer continuing into the pseudoparenchymatous cells of the erumpent stroma. Asci 8-spored, clavate, (71)–84–114(–142) × (8)–11–17(–22) µm (av. = 99 × 14 µm), tapering to a long thin stalk. Ascospores aggregate in the upper third of the ascii, hyaline, guttulate, fusoid with rounded ends, straight to slightly curved, 1(–3)-septate, not or slightly constricted at the septum, (24)–27–35(–43) × (4.5)–5.5–7.5(–9.5) µm (av. = 31 × 6.5 µm). Macroconidiophores consisting of a stipe, a suite of penicillate arranged fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 32–146 × 3.5–7.5 µm; stipe extension septate, straight to flexuous 35–217 µm long, 1.5–3.5 µm wide at the apical septum, terminating in a sphaeropedunculate vesicle, (4.5)–5–8.5(–12) µm diam; lateral stipe extensions (90° to main axis) abundant, septate, straight to flexuous 21–143 µm long, terminating in a sphaeropedunculate vesicle, 1–4 µm diam. Conidiogenous apparatus 32–187 µm long, 23–126 µm wide; primary branches aseptate to 1-septate, 13–53 × 3.5–6 µm; secondary branches aseptate, 11–28 × 3–5.5 µm; tertiary branches aseptate, 8.5–21 × 3–5 µm; additional branches (–5), aseptate, 6–11.5 × 2–4.5 µm; each terminal branch producing 2–4 phialides; phialides doliiiform

to reniform, hyaline, aseptate, 7–13.5 × 2–4.5 µm, apex with minute periclinal thickening and inconspicuous collarette. Macroconidia cylindrical, rounded at both ends, straight, (33)–36–44(–50) × (2.5)–3.5–4 µm, (av. = 40 × 3.5 µm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. Mega- and microconidia not observed.

**Culture characteristics:** Colonies forming abundant white to saffron aerial mycelium on MEA at 25 °C after 7 d, profuse sporulation, feathery, irregular margins, reverse capucine buff (13f) to russet (13'k). Chlamydospores common throughout the medium forming microsclerotia. **Growth characteristics:** optimal growth temperature 25 °C, no growth at 5 °C and 35 °C, after 7 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C and 30 °C reached 9.3 mm, 28.9 mm, 43.3 mm, 68.7 mm and 57.2 mm, respectively.

**Additional material examined:** **China:** FuJian Province: Fuzhou City, MinHou County, BaiSha Town: from soil collected in a *Eucalyptus* plantation, 12 Dec. 2014, S.F. Chen (PREM 61947, culture CMW 47489 = CERC 7115 = CBS 142889); FuZhou City, CangShan District: from soil collected in the campus of Fujian Agriculture and Forestry University (FAFU), 14 Dec. 2014, S.F. Chen (PREM 61949, culture CMW 47497 = CERC 7127 = CBS 142891).

**Notes:** *Calonectria pseudoturangicola* is phylogenetically closely related to *Ca. hongkongensis* and *Ca. turangicola*, but the macroconidia of *Ca. pseudoturangicola* (av. = 40 × 3.5 µm) are shorter and narrower than those of *Ca. hongkongensis* (av. = 46.5 × 4 µm) and *Ca. turangicola* (av. = 44 × 4 µm) (Crous et al. 2004, Lombard et al. 2015).

**Calonectria pseudoyunnanensis** J.Q. Li, Q.L. Liu & S.F. Chen, sp. nov.


MycoBank MB821636  
(Fig. 7)

**Etymology:** From the close resemblance to *Calonectria yunnanensis*.


**Diagnosis:** *Calonectria pseudoyunnanensis* can be distinguished from the phylogenetically closely related *Ca. asiatica*, *Ca. colombiensis*, and *Ca. yunnanensis* by the size of macroconidia and the shape of vesicles.

**Type:** **China:** YunNan Province: PuEr Region, JingGu County, WeiYuan Town, from soil collected in a *Eucalyptus* plantation, 16 Nov. 2014, S.F. Chen & J.Q. Li (PREM 61950 – holotype; CMW 47655 = CERC 5376 = CBS 142892 – ex-type cultures).

**Description:** Sexual morph unknown. Macroconidiophores consisting of a stipe, a suite of penicillate arranged fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 38–89 × 5–8 µm; stipe extension septate, straight to flexuous 22–94 µm long, 1.5–2.5 µm wide at the apical septum, terminating in ellipsoidal, obpyriform to sphaeropedunculate vesicles, (2.5)–3.5–5.0 µm diam; lateral stipe extensions (90° to main axis) abundant, septate, straight to flexuous 18–64 µm long, terminating in a obpyriform to



**Fig. 6.** *Calonectria pseudoturangicola*. **A.** Perithecium. **B.** Vertical section through a perithecium. **C.** Cells around ostiolar region of perithecium. **D.** Section through lateral perithecial wall. **E–F.** Ascii. **G.** Ascospores. **H–I.** Macroconidiophore. **J–K.** Sphaeropedunculate vesicles. **L–M.** Conidiogenous apparatus with conidiophore branches and doliiform to reniform phialides. **N.** Macroconidia. Bars: A = 200 µm; B = 50 µm; C–F and H–I = 20 µm; G and J–N = 10 µm.



**Fig. 7.** *Calonectria pseudoyunnanensis*. **A–B.** Macroconidiophore. **C–F.** Ellipsoidal, obovoidal to sphaeropedunculate vesicles. **G–H.** Conidiogenous apparatus with conidiophore branches and doliiform to reniform phialides. **I–J.** Macroconidia. Bars: A–B = 20 µm; G–J = 10 µm; C–F = 5 µm.

sphaeropedunculate vesicle, 1–3 µm diam. *Conidiogenous apparatus* 28–87 µm long, 32–83 µm wide; primary branches aseptate to 1-septate, 16–42 × 3.5–6.5 µm; secondary branches aseptate, 11–19 × 3.5–5.5 µm; tertiary branches aseptate, 7–13 × 3–5 µm; each terminal branch producing 2–4 phialides; phialides doliform to reniform, hyaline, aseptate, 6–15 × 3–5 µm; apex with minute periclinal thickening and inconspicuous collarette. *Macroconidia* cylindrical, rounded at both ends, straight, (40–)44–50(–55) × (4–)4.5–5.5(–6) µm (av. = 47.5 × 5 µm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not observed.

**Culture characteristics:** Colonies forming abundant white aerial mycelium on MEA at 25 °C after 7 d, moderate sporulation, feathery, irregular margins, reverse pale yellow-orange (15f) to sienna (8). *Chlamydospores* common throughout the medium forming microsclerotia. **Growth characteristics,** optimal growth temperature 25 °C, no growth at 5 °C and 35 °C, after 7 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C and 30 °C reached 6.4 mm, 30.4 mm, 53.7 mm, 78.4 mm and 55.3 mm, respectively.

**Additional material examined:** **China:** YunNan Province: PuEr Region, JingGu County, WeiYuan Town, from soil collected in a *Eucalyptus* plantation, 16 Nov. 2014, S.F. Chen & J.Q. Li (PREM 61951, CMW 47656 = CERC 5377 = CBS 142893; PREM 61952, culture CMW 47657 = CERC 5378 = CBS 142894).

**Notes:** *Calonectria pseudoyunnanensis* is most closely related to *Ca. asiatica*, *Ca. colombiensis*, and *Ca. yunnanensis*. It can be distinguished from these three species by the average size of the macroconidia. Those of *Ca. pseudoyunnanensis* (av. = 47.5 × 5 µm) are longer and broader than those of *Ca. yunnanensis* (av. = 43 × 4.5 µm), but shorter than those of *Ca. asiatica* (av. = 53 × 5 µm) and *Ca. colombiensis* (av. = 53 × 4.5 µm) (Crous et al. 2004). Furthermore, the vesicle shape of *Ca. pseudoyunnanensis* (ellipsoidal, obpyriform to sphaeropedunculate) is different to those of *Ca. asiatica* (sphaeropedunculate) and *Ca. colombiensis* (sphaeropedunculate) (Crous et al. 2004, Lombard et al. 2010c).

**Calonectria yunnanensis** J.Q. Li, Q.L. Liu & S.F. Chen, sp. nov.  
Mycobank MB821637  
(Fig. 8)

**Etymology:** From YunNan Province, China, where this fungus was first collected.

**Diagnosis:** *Calonectria yunnanensis* can be distinguished from the phylogenetically closely related *Ca. asiatica*, *Ca. colombiensis*, and *Ca. pseudoyunnanensis* by the size of macroconidia and ascospores.


**Type:** **China:** YunNan Province: PuEr Region, JingGu County, ZhengXing Town, from soil collected in a *Eucalyptus* plantation, 16 Nov. 2014, S.F. Chen & J.Q. Li (PREM 61955 – holotype; CMW 47644 = CERC 5339 = CBS 142897 – ex-type cultures).

**Description:** *Perithecia* solitary or in groups of up to five, orange, becoming orange-brown with age; in section, apex and body orange, base red-brown, subglobose to ovoid, 303–511 µm high, 322–567 µm diam, body turning red, and base dark red-brown in 3 % KOH; perithecial walls rough consisting of two thick-walled layers: outside layer of *textura globulosa*, 24–72 µm wide, becoming more compressed towards inner layer of *textura angularis*, 10–22 µm wide, becoming thin-walled and hyaline towards the centre; outer cells 19–37 × 12–21 µm, inner cells 14–39 × 3–11 µm; perithecial base up to 260 µm wide, consisting of dark red, angular cells merging with an erumpent stroma, cells of the outer wall layer continuing into the pseudoparenchymatous cells of the erumpent stroma. Asci 8-spored, clavate, (84–)97–133(–163) × (10–)15–21(–27) µm (av. = 115 × 18 µm), tapering to a long thin stalk. Ascospores aggregate in the upper third of the asci, hyaline, guttulate, fusoid with rounded ends, straight to slightly curved, 1(–3)-septate, not or slightly constricted at the septum, (28–)31–41(–55) × (5–)5.5–6.5(–8) µm (av. = 36 × 6 µm). *Macroconidiophores* consisting of a stipe, a suite of penicillate arranged fertile branches, a stipe extension, and a terminal vesicle; stipe septate, hyaline, smooth, 43–230 × 2.5–7 µm; stipe extension septate, straight to flexuous 25–102 µm long, 1.5–3.5 µm wide at the apical septum, terminating in a sphaeropedunculate vesicle, (2–)2.5–3.5(–4.5) µm diam; lateral stipe extensions (90° to main axis) abundant, septate, straight to flexuous 25–69 µm long, terminating in a sphaeropedunculate vesicle, 1–4 µm diam. *Conidiogenous apparatus* 20–130 µm long, 23–135 µm wide; primary branches aseptate to 1-septate, 13–49 × 3–6.5 µm; secondary branches aseptate, 12–17 × 3–5 µm; tertiary branches aseptate, 4–13 × 1.5–4 µm; each terminal branch producing 2–4 phialides; phialides doliform to reniform, hyaline, aseptate, 6–16 × 2.5–4.5 µm, apex with minute periclinal thickening and inconspicuous collarette. *Macroconidia* cylindrical, rounded at both ends, straight, (36–)39–47(–52) × (4–)4.5–5(–5.5) µm, (av. = 43 × 4.5 µm), 1-septate, lacking a visible abscission scar, held in parallel cylindrical clusters by colourless slime. *Mega-* and *microconidia* not observed.

**Culture characteristics:** Colonies forming abundant white to white-buff aerial mycelium on MEA at 25 °C after 7 d, profuse sporulation, feathery, irregular margins, reverse salmon (13'd) to sienna (8). *Chlamydospores* common throughout the medium forming microsclerotia. **Growth characteristics,** optimal growth temperature 25 °C, no growth at 5 °C and 35 °C, after 7 d, colonies at 10 °C, 15 °C, 20 °C, 25 °C and 30 °C reached 7.3 mm, 33.0 mm, 53.9 mm, 76.4 mm and 53.9 mm, respectively.

**Additional material examined:** **China:** YunNan Province: PuEr Region, JingGu County, ZhengXing Town, from soil collected in a *Eucalyptus* plantation, 16 Nov. 2014, S.F. Chen & J.Q. Li (PREM 61953, culture CMW 47642 = CERC 5337 = CBS 142895; PREM 61954, culture CMW 47643 = CERC 5338 = CBS 142896).

**Notes:** *Calonectria yunnanensis* is closely related to *Ca. asiatica*, *Ca. colombiensis* and *Ca. pseudoyunnanensis*. It can be distinguished from these three species by the



**Fig. 8.** *Calonectria yunnanensis*. **A.** Perithecium. **B.** Vertical section through a perithecium. **C.** Cells around ostiolar region of perithecium. **D.** Section through lateral perithecial wall. **E–F.** Asci. **G.** Ascospores. **H–I.** Macroconidiophore. **J–K.** Sphaeropedunculate vesicles. **L–M.** Conidiogenous apparatus with conidiophore branches and doliiiform to reniform phialides. **N.** Macroconidia. Bars: A = 200 µm; B = 100 µm; E = 50 µm; C–D, F and H–I = 20 µm; G and J–N = 10 µm.

average size of the macroconidia. The macroconidia of *Ca. yunnanensis* (av. =  $43 \times 4.5 \mu\text{m}$ ) are shorter than those of *Ca. asiatica* (av. =  $53 \times 5 \mu\text{m}$ ), *Ca. colombiensis* (av. =  $53 \times 4.5 \mu\text{m}$ ) and *Ca. pseudoyunnanensis* (av. =  $47.5 \times 5 \mu\text{m}$ ) (Crous et al. 2004). The ascospores of *Ca. yunnanensis* (av. =  $36 \times 6 \mu\text{m}$ ) are slightly longer than those of *Ca. asiatica* (av. =  $33 \times 6 \mu\text{m}$ ) and *Ca. colombiensis* (av. =  $33 \times 5 \mu\text{m}$ ) (Crous et al. 2004).

## DISCUSSION

Results of this study revealed 13 species of *Calonectria* from infected *Eucalyptus* tissues collected in plantations and nurseries, or baited soil samples from *Eucalyptus* plantations, and a naturally forested area in South China. These species include *Ca. arbusta*, *Ca. asiatica*, *Ca. chinensis*, *Ca. eucalypti*, *Ca. hongkongensis*, *Ca. mossambicensis*, *Ca. pentaseptata*, and six previously undescribed taxa (*Ca. aciculata*, *Ca. honghensis*, *Ca. lantauensis*, *Ca. pseudoturangicola*, *Ca. pseudoyunnanensis*, and *Ca. yunnanensis*). The six novel species were strongly supported by DNA sequence data and morphological observations. Five of the 13 species, including *Ca. aciculata*, *Ca. eucalypti*, *Ca. honghensis*, *Ca. mossambicensis*, and *Ca. pentaseptata*, resided in the Prolate Group and eight in the Sphaero-Naviculate Group. With the exception of the newly described species, this is the first report of *Ca. asiatica*, *Ca. eucalypti*, and *Ca. mossambicensis* from China. *Calonectria chinensis* and *Ca. lantauensis* were isolated only from soil in natural forested areas, while 11 species were all collected from *Eucalyptus* plantations or nurseries.

*Calonectria pentaseptata*, identified in this study, resides in the *Ca. reteaudii* species complex and was widely distributed in different regions causing disease on *Eucalyptus* in plantations and nurseries in South China. Amongst the 115 *Calonectria* isolates collected in this study, approximately half (57) were identified as *Ca. pentaseptata*, and this fungus occurred at six different sites in the GuangDong and GuangXi Provinces. This is consistent with previous studies showing that *Ca. pentaseptata* is widely distributed in *Eucalyptus* plantations and nurseries in South China (Lombard et al. 2015). The *Ca. reteaudii* complex, which includes species that are well-known causal agents of *Calonectria* Leaf Blight (CLB) of *Eucalyptus* (Crous 2002, Rodas et al. 2005, Lombard et al. 2010b). *Calonectria pentaseptata* is the fourth species in the *Ca. reteaudii* species complex to have been found in China; the other three include *Ca. crousiiana*, *Ca. microconidialis* and *Ca. pseudoreteaudii*. Pathogenicity tests have shown that all four of these species cause rot on inoculated *Eucalyptus* leaves (Chen et al. 2011c, Li et al. 2014a, b). Overall, the results of this study support the view (Lombard et al. 2015) that *Ca. pentaseptata* is an important *Eucalyptus* pathogen both in plantations and nurseries in China.

*Calonectria mossambicensis* is the fourth species in the *Ca. candelabrum* complex to have been reported from China together with *Ca. pauciramosa*, *Ca. seminaria* and *Ca. tetraramosa*. All four species were isolated from diseased seedlings in *Eucalyptus* nurseries (Lombard et al. 2010d, 2015). Species in the *Ca. candelabrum* complex include

some important nursery pathogens (Crous 2002, Lombard et al. 2010b, d, Guarnaccia et al. 2014, Alfenas et al. 2015). Inoculation studies have also shown that *Ca. pauciramosa*, *Ca. seminaria* and *Ca. tetraramosa* are differentially pathogenic to *Eucalyptus* clones (Chen et al. 2011c, Li et al. 2014a, b). *Calonectria mossambicensis* was originally described from diseased cuttings of *E. grandis* × *E. camaldulensis* clones in Mozambique (Crous et al. 2013) and it is likely to be a *Eucalyptus* nursery pathogen in China, since this fungus causes rot on *Eucalyptus* cutting rot in Mozambique.

Three species residing in the *Ca. colhounii* complex were identified in this study. They include *Ca. eucalypti* and the newly described *Ca. aciculata* and *Ca. honghensis*. Species in the *Ca. colhounii* complex are characterized by bright yellow perithecia (Crous 2002, Lombard et al. 2010c, Chen et al. 2011c, Xu et al. 2012). *Calonectria aciculata* and *Ca. honghensis* are closely related to *Ca. colhounii* and these three species can easily be distinguished from each other based on phylogenetic inference, as well as by their macroconidial dimensions. Other *Calonectria* species known in China and that reside in the *Ca. colhounii* complex include *Ca. fujianensis*, *Ca. nymphaeae*, and *Ca. pseudocolhounii* (Chen et al. 2011c, Xu et al. 2012). Other than *Ca. honghensis* isolated from soil collected in a *Eucalyptus* plantation, and *Ca. nymphaeae* from diseased leaves of *Nymphaea tetragona* (Xu et al. 2012), the remaining four species in the *Ca. colhounii* complex were all isolated from diseased *Eucalyptus* leaves in commercial plantations (Chen et al. 2011c). Inoculation studies have shown that *Ca. crousiiana*, *Ca. fujianensis* and *Ca. pseudocolhounii* are all pathogenic to inoculated *Eucalyptus* clones (Chen et al. 2011c, Li et al. 2014a, b).

Four new species in the Sphaero-Naviculate Group reside in the *Ca. kyotensis* complex, *Ca. lantauensis*, *Ca. pseudoturangicola*, *Ca. pseudoyunnanensis*, and *Ca. yunnanensis*. *Calonectria pseudoyunnanensis* and *Ca. yunnanensis* are sister species based on phylogenetic inference but they can easily be distinguished by DNA sequence comparisons of the *his3* and *tub2* gene regions, and vesicle shape differences. *Calonectria pseudoturangicola* appears as a sister species to *Ca. turangicola* but can be distinguished based on DNA sequence differences in the *tef1*, *cmdA* and *tub2* gene regions, and macroconidial dimensions (Lombard et al. 2015). *Calonectria lantauensis* formed a basal clade in the *Ca. kyotensis* species complex, and lateral stipe extensions were absent in this species making it readily distinguishable from other species in the *Ca. kyotensis* species complex (Crous et al. 2004, Lombard et al. 2010c, 2015).

The remaining four known species (*Ca. arbusta*, *Ca. asiatica*, *Ca. chinensis*, and *Ca. hongkongensis*) found in this study reside in the *Ca. kyotensis* complex. To date, 19 species in the *Ca. kyotensis* complex have been found in China and the only other species in the complex, *Ca. asiatica*, was first described from Thailand (Crous et al. 2004, Lombard et al. 2015). These 19 species were all isolated exclusively from soil (Lombard et al. 2015) and the results of this study suggest that many more species in this complex have yet to be discovered from soil in China.

Overall, the results of this study revealed 37 species

of *Calonectria* from China. Other than *Ca. asiatica*, *Ca. eucalypti*, *Ca. mossambicensis*, *Ca. pauciramosa*, and *Ca. pentaseptata*, all of these species were first discovered in this country (Crous *et al.* 2004, Lombard *et al.* 2010d, 2015, Chen *et al.* 2011c, Xu *et al.* 2012). The results highlight the significant impact that DNA sequence comparisons have had in revealing new species of filamentous fungi, including species of *Calonectria* (Lombard *et al.* 2010c, 2015, 2016, Crous *et al.* 2015a).

With the exception of *Ca. nymphaeae* isolated from diseased leaves of *N. tetragona* (Xu *et al.* 2012), and *Ca. lantauensis* from a naturally forested area in Hong Kong, all of the other 35 species found in China were from *Eucalyptus* plantations or nurseries. This appears to be an environment surprisingly rich in species of *Calonectria*, although future sampling in China should be expanded to include other environments. Inoculation tests conducted in previous studies have shown that 15 species of *Calonectria* found in China are pathogenic to several *Eucalyptus* clones (Chen *et al.* 2011c, Li *et al.* 2014a, b). Future work should include a more comprehensive understanding of the species diversity, distribution, pathogenicity and population biology of *Calonectria* in China. This will contribute to the development of integrated management strategies for the diseases caused by these fungi in *Eucalyptus* plantations and nurseries.

## ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China (NSFC) (Project Numbers 31622019, 31400546). The authors acknowledge members of Tree Protection and Cooperation Programme (TPCP). The authors thank GuoQing Li, FeiFei Liu, and MiRu Zhang for their assistance in collecting samples.

## REFERENCES

- Alfenas RF, Lombard L, Pereira OL, Alfenas AC, Crous PW (2015) Diversity and potential impact of *Calonectria* species in *Eucalyptus* plantations in Brazil. *Studies in Mycology* **80**: 89–130.
- Brown BN, Ferreira FA (2000) *Disease during Propagation of Eucalypts*. Canberra: CSIRO publishing.
- Burgess TI, Andjic V, Hardy GS, Dell B, Xu D (2006) First report of *Phaeophleospora destructans* in China. *Journal of Tropical Forest Science* **18**: 144–146.
- Burgess TI, Barber PA, Sufaati S, Xu D, Hardy GS, *et al.* (2007) *Mycosphaerella* spp. on *Eucalyptus* in Asia: new species, new hosts and new records. *Fungal Diversity* **24**: 135–157.
- Chen SF, Gryzenhout M, Roux J, Xie YJ, Wingfield MJ, *et al.* (2010) Identification and pathogenicity of *Chrysoporthe cubensis* on *Eucalyptus* and *Syzygium* spp. in South China. *Plant Disease* **94**: 1143–1150.
- Chen SF, Barnes I, Chungu D, Roux J, Wingfield MJ, *et al.* (2011a) High population diversity and increasing importance of the *Eucalyptus* stem canker pathogen, *Teratosphaeria zuluensis*, in South China. *Australasian Plant Pathology* **40**: 407–415.
- Chen SF, Gryzenhout M, Roux J, Xie YJ, Wingfield MJ, *et al.* (2011b) Novel species of *Celoporthe* from *Eucalyptus* and *Syzygium* trees in China and Indonesia. *Mycologia* **103**: 1384–1410.
- Chen SF, Lombard L, Roux J, Xie YJ, Wingfield MJ, *et al.* (2011c) Novel species of *Calonectria* associated with *Eucalyptus* leaf blight in Southeast China. *Persoonia* **26**: 1–12.
- Chen SF, Pavlic D, Roux J, Slippers B, Xie YJ, *et al.* (2011d) Characterization of *Botryosphaeriaceae* from plantation-grown *Eucalyptus* species in South China. *Plant Pathology* **60**: 739–751.
- Chen SF, Van Wyk M, Roux J, Wingfield MJ, Xie YJ, *et al.* (2013) Taxonomy and pathogenicity of *Ceratocystis* species on *Eucalyptus* trees in South China, including *C. chinæucensis* sp. nov. *Fungal Diversity* **58**: 267–279.
- Chen SX, Chen XF (2013) Technical problems and thinking on eucalypt plantation management in China. *Eucalypt Science & Technology* **30**: 52–59. [In Chinese]
- Cortinas MN, Burgess T, Dell B, Xu D, Crous PW, *et al.* (2006) First record of *Colletogloeopsis zuluense* comb. nov., causing a stem canker of *Eucalyptus* in China. *Mycological Research* **110**: 229–236.
- Crous PW (2002) *Taxonomy and Pathology of Cyphomycetes (Calonectria) and allied genera*. St Paul, MN: American Phytopathological Society Press.
- Crous PW, Groenewald JZ, Risède JM, Simoneau P, Hywel-Jones NL (2004) *Calonectria* species and their *Cyphomycetes* anamorphs: species with sphaeropedunculate vesicles. *Studies in Mycology* **50**: 415–430.
- Crous PW, Phillips AJL, Wingfield MJ (1991) The genera *Cyphomycetes* and *Cyphomycetidae* in South Africa, with special reference to forest nurseries. *South African Forestry Journal* **157**: 69–85.
- Crous PW, Shivas RG, Wingfield MJ, Summerell BA, Rossman AY, *et al.* (2012) Fungal Planet description sheets: 128–153. *Persoonia* **29**: 146–201.
- Crous PW, Wingfield MJ, Guarro J, Cheewangkoon R, Van der Bank M, *et al.* (2013) Fungal Planet description sheets: 154–213. *Persoonia* **31**: 188–296.
- Crous PW, Hawksworth DL, Wingfield MJ (2015a) Identifying and naming plant-pathogenic fungi: past, present, and future. *Annual Review of Phytopathology* **53**: 247–267.
- Crous PW, Wingfield MJ, Le Roux JJ, Richardson DM, Strasberg D, *et al.* (2015b) Fungal Planet description sheets: 371–399. *Persoonia* **35**: 264–327.
- Cunningham CW (1997) Can three incongruence tests predict when data should be combined? *Molecular Biology and Evolution* **14**: 733–740.
- Dettman JR, Jacobson DJ, Taylor JW (2003) A multilocus genealogical approach to phylogenetic species recognition in the model eukaryote *Neurospora*. *Evolution* **57**: 2703–2720.
- Geurber JC, Correll JC (2001) Characterization of *Glomerella acutata*, the teleomorph of *Colletotrichum acutatum*. *Mycologia* **93**: 216–229.
- Gilligan CA (1983) Modeling of soilborne pathogens. *Annual Review of Phytopathology* **21**: 45–64.
- Guarnaccia V, Aiello D, Polizzi G, Perrone G, Stea G, *et al.* (2014) Emergence of prochloraz-resistant populations of *Calonectria pauciramosa* and *Calonectria polizzii* in ornamental nurseries of southern Italy. *Plant Disease* **98**: 344–350.
- Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology* **52**: 696–704.
- Hillis DM, Bull JJ (1993) An empirical test of bootstrapping as a

- method for assessing confidence in phylogenetic analysis. *Systematic Biology* **42**: 182–192.
- Hwang SC, Ko WH (1976) Biology of conidia, ascospores, and microsclerotia of *Calonectria crotalariae* in soil. *Phytopathology* **66**: 51–54.
- Katho K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**: 772–780.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, et al. (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**: 1647–1649.
- Li GQ, Chen SF, Wu ZH, Zhou XD, Xie YJ (2014a) Preliminary analyses on diversity and pathogenicity of *Calonectria* spp. on *Eucalyptus* in China. *Chinese Journal of Tropical Crops* **35**: 1183–1191. [In Chinese]
- Li GQ, Li JQ, Liu FF, Li TH, Chen SF (2014b) Preliminary analyses on pathogenicity of twelve *Calonectria* spp. on ten *Eucalyptus* clones in China. *Eucalypt Science & Technology* **31**: 1–7. [In Chinese]
- Lombard L, Crous PW, Wingfield BD, Wingfield MJ (2010a) Species concepts in *Calonectria* (*Cylindrocladium*). *Studies in Mycology* **66**: 1–14.
- Lombard L, Crous PW, Wingfield BD, Wingfield MJ (2010b) Multigene phylogeny and mating tests reveal three cryptic species related to *Calonectria pauciramosa*. *Studies in Mycology* **66**: 15–30.
- Lombard L, Crous PW, Wingfield BD, Wingfield MJ (2010c) Phylogeny and systematics of the genus *Calonectria*. *Studies in Mycology* **66**: 31–69.
- Lombard L, Zhou XD, Crous PW, Wingfield BD, Wingfield MJ (2010d) *Calonectria* species associated with cutting rot of *Eucalyptus Persoonia* **24**: 1–11.
- Lombard L, Chen SF, Mou X, Zhou XD, Crous PW, et al. (2015) New species, hyper-diversity and potential importance of *Calonectria* spp. from *Eucalyptus* in South China. *Studies in Mycology* **80**: 151–188.
- Lombard L, Wingfield MJ, Alfenas AC, Crous PW (2016) The forgotten *Calonectria* collection: pouring old wine into new bags. *Studies in Mycology* **85**: 159–198.
- Marin-Felix Y, Groenewald JZ, Cai L, Chen Q, Marincowitz S, et al. (2017) Genera of phytopathogenic fungi: GOPHY 1. *Studies in Mycology* **86**: 99–216.
- Nirenberg HI (1981) A simplified method for identifying *Fusarium* spp. occurring on wheat. *Canadian Journal of Botany* **59**: 1599–1609.
- Ogden TH, Rosenberg MS (2007) How should gaps be treated in parsimony? A comparison of approaches using simulation. *Molecular Phylogenetics and Evolution* **42**: 817–826.
- Posada D (2008) jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* **25**: 1253–1256.
- Rayner RW (1970) *A Mycological Colour Chart*. Kew: Commonwealth Mycological Institute.
- Rodas CA, Lombard L, Gryzenhout M, Slippers B, Wingfield MJ (2005) *Cylindrocladium* blight of *Eucalyptus grandis* in Colombia. *Australasian Plant Pathology* **34**: 143–149.
- Schoch CL, Crous PW, Wingfield MJ, Wingfield BD (2000) Phylogeny of *Calonectria* and selected hypocrealean genera with cylindrical macroconidia. *Studies in Mycology* **45**: 45–62.
- Swofford DL (2003) *PAUP\**: phylogenetic analysis using parsimony (\*and other methods). Version 4. Sunderland, MA: Sinauer Associates.
- Tamura K, Stecher G, Peterson D, Filipski A, Sudhir Kumar (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* **30**: 2725–2729.
- Thies WG, Patton RF (1970) The biology of *Cylindrocladium scoparium* in Wisconsin forest tree nurseries. *Phytopathology* **60**: 1662–1668.
- Van Burik JAH, Schreckhise RW, White TC, Bowden RA, Myerson D (1988) Comparison of six extraction techniques for isolation of DNA from filamentous fungi. *Medical Mycology* **36**: 299–303.
- Xu JJ, Qin SY, Hao YY, Ren J, Tan P, et al. (2012) A new species of *Calonectria* causing leaf disease of water lily in China. *Mycotaxon* **122**: 177–185.
- Zhou XD, De Beer ZW, Xie YJ, Pegg GS, Wingfield MJ (2007) DNA-based identification of *Quambalaria pitereka* causing severe leaf blight of *Corymbia citriodora* in China. *Fungal Diversity* **25**: 245–254.
- Zhou XD, Xie YJ, Chen SF, Wingfield MJ (2008) Diseases of eucalypt plantations in China: challenges and opportunities. *Fungal Diversity* **32**: 1–7.

**Supplementary Table 1**

 Supplementary Table 1. The species of *Calonectria* previously reported associated with *Eucalyptus* in China.

Species No.	Species	Phylogenetic Group	Substrate	Locality	Symptoms/Pathogenicity	References
1	<i>Calonectria aconidialis</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	HaiNan	NA <sup>a</sup> /NA	Lombard <i>et al.</i> 2015
2	<i>Ca. arbusta</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi	NA/NA	Lombard <i>et al.</i> 2015
3	<i>Ca. cerciana</i>	Prolate	<i>Eucalyptus urophylla</i> × <i>Eucalyptus grandis</i> cutting stem in nursery	GuangDong	Stem rot/leaf rot of seedling	Lombard <i>et al.</i> 2010d; Li <i>et al.</i> 2014a, b
4	<i>Ca. chinensis</i>	Sphaero-Naviculate	Soil in natural forestry and <i>Eucalyptus</i> plantation	Hong Kong, GuangDong, HaiNan	NA/leaf rot of seedling	Crous <i>et al.</i> 2004; Li <i>et al.</i> 2014b; Lombard <i>et al.</i> 2015
5	<i>Ca. crousiana</i>	Prolate	<i>E. grandis</i> leaf in plantation	Fujian	CLB <sup>b</sup> /leaf rot of seedling	Chen <i>et al.</i> 2011c; Li <i>et al.</i> 2014a
6	<i>Ca. expansa</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangDong, GuangXi	NA/NA	Lombard <i>et al.</i> 2015
7	<i>Ca. foliicola</i>	Prolate	<i>E. urophylla</i> × <i>E. grandis</i> leaf in plantation	GuangXi	CLB/NA	Lombard <i>et al.</i> 2015
8	<i>Ca. fujianensis</i>	Prolate	<i>E. grandis</i> leaf in plantation	Fujian	CLB/leaf rot of seedling	Chen <i>et al.</i> 2011c; Li <i>et al.</i> 2014a
9	<i>Ca. guangxiensis</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi	NA/NA	Lombard <i>et al.</i> 2015
10	<i>Ca. hainanensis</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	HaiNan	NA/NA	Lombard <i>et al.</i> 2015
11	<i>Ca. hongkongensis</i>	Sphaero-Naviculate	Soil in natural forestry and <i>Eucalyptus</i> plantation	Hong Kong, GuangDong, GuangXi, HaiNan	NA/leaf rot of seedling	Crous <i>et al.</i> 2004; Li <i>et al.</i> 2014b
12	<i>Ca. lateralis</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi	NA/NA	Lombard <i>et al.</i> 2015
13	<i>Ca. magnispora</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi	NA/NA	Lombard <i>et al.</i> 2015
14	<i>Ca. microcondialis</i>	Prolate	<i>E. urophylla</i> × <i>E. grandis</i> seedling leaf in nursery	GuangDong	Leaf rot of seedling/leaf rot of seedling	Li <i>et al.</i> 2014b; Lombard <i>et al.</i> 2015
15	<i>Ca. papillata</i>	Prolate	Soil in <i>Eucalyptus</i> plantation	GuangDong, GuangXi	NA/leaf rot of seedling	Li <i>et al.</i> 2014b; Lombard <i>et al.</i> 2015
16	<i>Ca. parakyotensis</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangDong, GuangXi	NA/leaf rot of seedling	Li <i>et al.</i> 2014b; Lombard <i>et al.</i> 2015
17	<i>Ca. pauciramosa</i>	Prolate	<i>Eucalyptus</i> leaf in plantation and <i>Eucalyptus</i> seedling stem in nursery	GuangDong, Fujian	CLB, stem rot/leaf rot of seedling	Chen <i>et al.</i> 2011c; Li <i>et al.</i> 2014a, b
18	<i>Ca. pentaseptata</i>	Prolate	<i>Eucalyptus</i> leaf and soil in <i>Eucalyptus</i> plantation and <i>Eucalyptus</i> seedling leaf in nursery	GuangDong, GuangXi, HaiNan,	CLB/leaf rot of seedling	Crous <i>et al.</i> 2012; Lombard <i>et al.</i> 2015
19	<i>Ca. pluriramosa</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi	NA/NA	Lombard <i>et al.</i> 2015
20	<i>Ca. pseudocoelounii</i>	Prolate	<i>E. dunnii</i> seedling leaf in nursery	Fujian	Leaf rot of seedling/leaf rot of seedling	Chen <i>et al.</i> 2011c; Li <i>et al.</i> 2014a
21	<i>Ca. pseudokyotensis</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi	NA/NA	Lombard <i>et al.</i> 2015
22	<i>Ca. pseudoreteaudi</i>	Prolate	<i>E. urophylla</i> × <i>E. grandis</i> cutting stem in nursery	GuangDong	Stem rot/leaf rot of seedling	Lombard <i>et al.</i> 2010d; Li <i>et al.</i> 2014a, b
23	<i>Ca. seminaria</i>	Prolate	<i>E. urophylla</i> × <i>E. grandis</i> seedling leaf in nursery	GuangDong	Leaf rot of seedling/leaf rot of seedling	Li <i>et al.</i> 2014b; Lombard <i>et al.</i> 2015
24	<i>Ca. sphaeropendunculata</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi	NA/NA	Lombard <i>et al.</i> 2015
25	<i>Ca. terrestris</i>	Prolate	<i>Eucalyptus</i> leaf and soil in <i>Eucalyptus</i> plantation	GuangDong, GuangXi	CLB/leaf rot of seedling	Li <i>et al.</i> 2014b; Lombard <i>et al.</i> 2015
26	<i>Ca. tetraramosa</i>	Prolate	<i>E. urophylla</i> × <i>E. grandis</i> seedling leaf in nursery	GuangDong	Leaf rot of seedling/leaf rot of seedling	Li <i>et al.</i> 2014b; Lombard <i>et al.</i> 2015
27	<i>Ca. turangicola</i>	Sphaero-Naviculate	Soil in <i>Eucalyptus</i> plantation	GuangXi, HaiNan	NA/NA	Lombard <i>et al.</i> 2015

<sup>a</sup> NA: No information is available.

<sup>b</sup> CLB: *Cylindrocladium* leaf blight.

**Supplementary Table 2**

Species	Isolate No. <sup>1</sup>	Substrate	Location	Collector	GenBank accession No. <sup>2</sup>			Reference
					<i>terf1</i>	<i>his3</i>	<i>cmaDA</i>	
<i>Ca. aconitidis</i>	<b>CBS 136086<sup>3</sup>;</b> CMW 35174; CERC 1850	Soil in <i>Eucalyptus</i> plantation	HaiNan, China	X. Mou & S.F. Chen	KJ462785	KJ463133	KJ463017	N/A <sup>d</sup> Lombard et al. (2015)
	<b>CBS 136091<sup>3</sup>;</b> CMW 35384; CERC 1886; CPC 23504	Soil in <i>Eucalyptus</i> plantation	HaiNan, China	X. Mou & S.F. Chen	KJ462786	KJ463134	N/A <sup>d</sup>	N/A Lombard et al. (2015)
	<b>CBS 136079<sup>3</sup>;</b> CMW 31370; CERC 1705; CPC 23482	Soil in <i>Eucalyptus</i> plantation	GuangXi, China	X. Zhou, G. Zhao & F.Han	KJ462787	KJ463135	KJ463018	KJ462904 Lombard et al. (2015)
	<b>CBS 136098<sup>3</sup>;</b> CMW 37981; CERC 1944; CPC 23519	Soil in <i>Eucalyptus</i> plantation	GuangXi, China	N/A <sup>5</sup>	KJ462788	KJ463136	KJ463019	N/A Lombard et al. (2015)
	<b>CBS 114073<sup>3</sup>;</b> CMW 23782; SFE 726; CPC 3900	Leaf litter	Prathet Thai, Thailand	N.L. Hywel- Jones	AY725705	AY725658	AY725741	AY725616 Crous et al. (2004); Lombard et al. (2010c)
	<b>CBS 112711<sup>3</sup>;</b> CPC 3898; SFE 744	Leaf litter	Prathet Thai, Thailand	N.L. Hywel- Jones	AY725702	AY725655	AY725738	AY725613 Crous et al. (2004); Lombard et al. (2010c)
<i>Ca. chinensis</i>	<b>CBS 114827<sup>3</sup>;</b> CMW 23674; CPC 4101	Soil	Hong Kong, China	E.C.Y. Liew	AY725710	AY725661	AY725747	AY725619 Crous et al. (2004); Lombard et al. (2010c)
	<b>CBS 112744<sup>3</sup>;</b> CMW 30986; CPC 4104	Soil	Hong Kong, China	E.C.Y. Liew	AY725709	AY725660	AY725746	AY725618 Crous et al. (2004); Lombard et al. (2010c)
	<b>CBS 293-79<sup>3</sup>;</b> CMW 30999	<i>Camellia sinensis</i>	Bandung, Indonesia	N/A	GQ267301	DQ190639	GQ267373	DQ190564 Crous (2002); Lombard et al. (2015)
<i>Ca. colombiensis</i>	<b>CBS 114704</b>	<i>Arachis pintoi</i>	Australia	D. Hutton	GQ267300	DQ190638	GQ267372	DQ190563 Crous (2002); Lombard et al. (2010c)
	<b>CBS 112220<sup>3</sup>;</b> CMW 23676; CPC 723	Soil under <i>Eucalyptus</i> <i>grandis</i> trees	La Selva, Colombia	M.J. Wingfield	AY725711	AY725662	AY725748	GQ267207 Crous et al. (2004); Lombard et al. (2010c)
	<b>CBS 112221<sup>3</sup>;</b> CMW 30985; CPC 724	Soil under <i>Eucalyptus</i> <i>grandis</i> trees	La Selva, Colombia	M.J. Wingfield	AY725712	AY725663	AY725749	AY725620 Crous et al. (2004); Lombard et al. (2010c)
<i>Ca. curvispora</i>	<b>CBS 116159<sup>3</sup>;</b> CMW 23693	Soil	Tamatave, Madagascar	P.W. Crous	GQ267302	AY725664	GQ267374	AF333394 Crous (2002); Lombard et al. (2010c)
	<b>CBS 125275<sup>3</sup>;</b> CMW 18444	<i>E. grandis</i> leaf	Aek Nauli, Sumatra Utara, Indonesia	M.J. Wingfield	GQ267338	GQ267267	GQ267430	GQ267218 Lombard et al. (2010c)

Supplementary Table 2. (Continued).

Species	Isolate No. <sup>1</sup>	Substrate	Location	Collector	GenBank accession No. <sup>2</sup>			Reference
					terf1	his3	cmda	
	CBS 125276; CMW 18445	<i>E. grandis</i> leaf	Aek Nauli, Sumatra Utara, Indonesia	M.J. Wingfield	GQ267339	GQ267268	GQ267431	GQ267219 Lombard et al. (2010c)
<i>Ca. expansa</i>	<b>CBS 136247</b> ; CMW 31392; CERC 1727	Soil in <i>Eucalyptus</i> plantation	GuangXi, China	X. Zhou, G. Zhao & F. Han	KJ462798	KJ463146	KJ462914	Lombard et al. (2015)
	CBS 136078; CMW 31441; CERC 1776	Soil in <i>Eucalyptus</i> plantation	FangChengGang, GuangXi, China	X. Zhou, G. Zhao & F. Han	KJ462797	KJ463145	KJ462928	KJ462913 Lombard et al. (2015)
<i>Ca. fujianensis</i>	<b>CBS 127201</b> ; CMW 27257	<i>E. grandis</i> leaf in plantation	Fujian, China	M.J. Wingfield	HQ285820	HQ285806	N/A	HQ285792 Chen et al. (2011c); Lombard et al. (2015), Chen et al. (2011c); Lombard et al. (2015)
	CBS 127200; CMW 27254	<i>E. grandis</i> leaf in plantation	Fujian, China	M.J. Wingfield	HQ285819	HQ285805	N/A	HQ285791 Chen et al. (2011c); Lombard et al. (2015)
<i>Ca. guangxiensis</i>	<b>CBS 136092</b> ; CMW 35409; CERC 1900; CPC 23506	Soil in <i>Eucalyptus</i> plantation	GuangXi, China	X. Mou & R. Chang	KJ462803	KJ463151	KJ462919	Lombard et al. (2015)
	CBS 136094; CMW 35411; CERC 1902; CPC 23507	Soil in <i>Eucalyptus</i> plantation	GuangXi, China	X. Mou & R. Chang	KJ462804	N/A	KJ463035	KJ462920 Lombard et al. (2015)
<i>Ca. hongkongensis</i>	<b>CBS 114828</b> ; CPC 4670	Soil	Hong Kong, China	M.J. Wingfield	AY725717	AY725667	AY725755	AY725622 Crouse et al. (2004); Lombard et al. (2010c)
	CBS 114711; CMW 30995; CPC 686	Soil	Hong Kong, China	M.J. Wingfield	AY725716	AY725666	AY725754	AY725621 Crouse et al. (2004); Lombard et al. (2010c)
<i>Ca. illicicola</i>	<b>CBS 190.50</b> ; CMW 30998; IMI 299389	<i>Solanum tuberosum</i>	Bogor Indonesia	K.B. Boedijin & J. Reitsma	AY725726	AY725676	AY725631	Crouse (2002); Lombard et al. (2015)
	CBS 112215; CPC 492	<i>Arachis hypogaea</i>	U.S.A.	Beute	AY725727	AY725684	AY725765	AY725639 Crouse et al. (2004); Lombard et al. (2015)
<i>Ca. indonesiae</i>	<b>CBS 112823</b> ; CMW 23683; CPC 4508	Soil	Warambunga, Indonesia	M.J. Wingfield	AY725718	AY725668	AY725756	AY725623 Crouse et al. (2004); Lombard et al. (2010c)
	CBS 112840; CPC 4554	<i>Syzygium aromaticum</i>	Warambunga, Indonesia	M.J. Wingfield	AY725720	AY725670	AY725758	AY725625 Crouse et al. (2004); Lombard et al. (2010c)
<i>Ca. lateralis</i>	<b>CBS 136629</b> ; CMW 31412; CERC 1747	Soil in <i>Eucalyptus</i> plantation	FangChengGang, GuangXi, China	X. Zhou, G. Zhao & F. Han	KJ462840	KJ463186	KJ463070	KJ462955 Lombard et al. (2015)
<i>Ca. microconidialis</i>	<b>CBS 136638</b> ; CMW 31487; CERC 1822	<i>E. urophylla</i> × <i>E. grandis</i> seedling leaf in nursery	SuiXi, ZhanJiang, GuangDong, China	G. Zhao	KJ462845	KJ463191	KJ463075	KJ462960 Lombard et al. (2015)

Supplementary Table 2. (Continued).

Species	Isolate No. <sup>1</sup>	Substrate	Location	Collector	GenBank accession No. <sup>2</sup>			Reference
					terf1	his3	cmda	
<i>Ca. monticola</i>	CBS 136633; CMW 31471; CERC 1806	<i>E. urophylla</i> × <i>E. grandis</i> seedling leaf in nursery	SuiXi, ZhanJiang, GuangDong, China	G. Zhao	KJ462842	KJ463188	KJ463072	KJ462957 Lombard et al. (2015)
	CBS 140645; CPC 28835	Soil	Chaing Mai, Thailand	P.W. Crous	KT964773	N/A	KT964771	KT964769 Crous et al. (2015b)
<i>Ca. mossambicensis</i>	CPC 28836	Soil	Chaing Mai, Thailand	P.W. Crous	KT964774	N/A	KT964772	KT964770 Crous et al. (2015b)
	CMW 36327	<i>E. grandis</i> × <i>E. camaldulensis</i> cutting in nursery	Bandula, Manica, Mozambique	J. Roux & S. Mause-Sitee	JX570718	JX570726	JX570722	N/A Crous et al. (2013)
<i>Ca. nymphaeae</i>	CMW 36329	<i>E. grandis</i> and <i>E. urophylla</i> cutting in nursery	Gurué, Zambézia	J. Roux & S. Mause-Sitee	JX570717	JX570725	JX570721	N/A Crous et al. (2013)
	CBS 131802; HGUP 100003	<i>Nymphaea</i> <i>tetragona</i> Georgi leaf	HuaXi, GuiYang, Guizhou, China	S.Y. Qin	KC555273	N/A	N/A	JN984864 Xu et al. (2012); Lombard et al. (2015)
<i>Ca. pacifica</i>	HGUP 100004	<i>N. tetragona</i> Georgi leaf	HuaXi, GuiYang, Guizhou, China	Y. Wang	KC555274	N/A	N/A	JN984865 Xu et al. (2012); Lombard et al. (2015)
	CBS 109063; CMW 16726; IMI 354528	<i>Araucaria</i> <i>heterophylla</i>	Hawaii, USA	M. Aragaki	AY725724	GQ267255	AY725762	GQ267213 Crous (2002); Lombard et al. (2015)
<i>Ca. paracollounii</i>	CBS 114038; CMW 30988; CPC 10717	<i>Ipomoea aquatica</i>	Auckland, New Zealand	C.F. Hill	GQ267320	AY725675	GQ267402	AY725630 Crous (2002); Lombard et al. (2015)
	CBS 114679; CPC 2445	Unknown	U.S.A.	A.Y. Rossman	KX784714	N/A	KX784582	KX784644 Lombard et al. (2016)
<i>Ca. parakyotensis</i>	CBS 114705; CPC 2423	Fruit of <i>Annona</i> <i>reticulata</i>	Australia	D. Hutton	KX784715	N/A	N/A	KX784645 Lombard et al. (2016)
	CBS 136085; CMW 35169; CERC 1845	Soil in <i>Eucalyptus</i> plantation	GuangDong, China	X. Mou & R. Chang	KJ462851	KJ463197	KJ463081	N/A Lombard et al. (2015)
<i>Ca. pauciramosa</i>	CBS 136095; CMW 35413; CERC 1904	Soil in <i>Eucalyptus</i> plantation	GuangXi, China	X. Mou & R. Chang	KJ462852	KJ463198	KJ463082	N/A Lombard et al. (2015)
	CMP 5683; CPC 971	<i>Eucalyptus</i> sp. in nursery.	Brazil	A.C. Alfenas	FJ918565	FJ918531	GQ267405	FJ918514 Crous (2002); Lombard et al. (2010c)
<i>Ca. pentaseptata</i>	CMW 30823; CPC 416	<i>E. grandis</i>	South Africa	P.W. Crous	FJ918566	FJ918532	GQ267404	FJ918515 Crous (2002); Lombard et al. (2010c)
	CBS 133349	<i>Eucalyptus</i> hybrid	Hanoi, Bavi, Vietnam	P.Q. Thu	JX855958	JX855946	N/A	JX855942 Crous et al. (2012); Lombard et al. (2015)

Supplementary Table 2. (Continued).

Species	Isolate No. <sup>1</sup>	Substrate	Location	Collector	GenBank accession No. <sup>2</sup>			Reference
					terf1	his3	cmdA	
<i>Ca. polizzi</i>	CBS 133351	<i>Macadamia</i> sp.	Hanoi, Bavi, Vietnam	P.Q. Thu	JX855960	JX855948	N/A	JX855944 Crous et al. (2012); Lombard et al. (2015)
	CBS 125270; CMW 7804; CPC 2681	<i>Callistemon</i> <i>citrinus</i>	Sicily, Italy	G. Polizzi	FJ972486	FJ972436	GQ267461	FJ972417 Lombard et al. (2010b, 2015)
	CBS 125271; CMW 10151; CPC 2771	<i>Arbutus unedo</i>	Sicily, Italy	G. Polizzi	FJ972487	FJ972437	GQ267462	FJ972418 Lombard et al. (2010b, 2015)
<i>Ca. pseudocolhounii</i>	<b>CBS 127195<sup>c</sup>;</b> CMW 27209	<i>E. dunnii</i> leaf in plantation	Fujian, China	M.J. Wingfield	HQ285816	HQ285802	N/A	HQ285788 Chen et al. (2011c); Lombard et al. (2015)
	CBS 127196; CMW 27213	<i>E. dunnii</i> leaf in plantation	Fujian, China	M.J. Wingfield	HQ285817	HQ285803	N/A	HQ285789 Chen et al. (2011c); Lombard et al. (2015)
<i>Ca. pseudokyotoensis</i>	<b>CBS 137332<sup>c</sup>;</b> CMW 31439; CERC 1774	Soil in <i>Eucalyptus</i> plantation	FangChengGang, GuangXi, China	X. Zhou, G. Zhao & F. Han	KJ462881	KJ463227	KJ463111	KJ462994 Lombard et al. (2015)
<i>Ca. pseudoreteaudi</i>	<b>CBS 123694<sup>c</sup>;</b> CMW 25310	<i>E. urophylla</i> × <i>E.</i> <i>grandis</i> cutting stem in nursery	SuiXi, ZhanJiang, GuangDong, China	M.J. Wingfield & X.D. Zhou	FJ918541	FJ918519	GQ267411	FJ918504 Lombard et al. (2010d, 2015)
	CBS 123696; CMW 25292	<i>E. urophylla</i> × <i>E.</i> <i>grandis</i> cutting stem in nursery	SuiXi, ZhanJiang, GuangDong, China	M.J. Wingfield & X.D. Zhou	FJ918542	FJ918520	GQ267410	FJ918505 Lombard et al. (2010d, 2015)
<i>Ca. reteaudi</i>	<b>CBS 112144<sup>c</sup>;</b> CMW 30984; CPC 3201	<i>E. camaldulensis</i> leaf	Vietnam	M.J. Dudzinski & P.Q. Thu	FJ918537	DQ190661	GQ267417	AF389833 Crous (2002); Lombard et al. (2010c)
	CBS 112143; CMW 16738; CPC 3200	<i>Eucalyptus</i> sp. leaf	Vietnam	M.J. Dudzinski & P.Q. Thu	FJ918536	DQ190660	GQ267418	GQ240642 Crous (2002); Lombard et al. (2010c)
<i>Ca. seminaria</i>	<b>CBS 136632<sup>c</sup>;</b> CMW 31450; CERC 1785; CPC 23488	<i>E. urophylla</i> × <i>E.</i> <i>grandis</i> seedling leaf in nursery	SuiXi, ZhanJiang, GuangDong, China	G. Zhao	KJ462885	KJ463231	KJ463115	KJ462998 Lombard et al. (2015)
	CBS 136639; CMW 31489; CERC 1824	<i>E. urophylla</i> × <i>E.</i> <i>grandis</i> seedling leaf in nursery	SuiXi, ZhanJiang, GuangDong, China	G. Zhao	KJ462886	KJ463232	KJ463116	KJ462999 Lombard et al. (2015)
<i>Ca. sumatrensis</i>	<b>CBS 112829<sup>c</sup>;</b> CMW 23698; CPC 4518	Soil under <i>Eucalyptus</i> tree	Northern Sumatra, Indonesia	M.J. Wingfield	AY725733	AY725696	AY725771	AY725649 Crous et al. (2004); Lombard et al. (2015)

Supplementary Table 2. (Continued).

Species	Isolate No. <sup>1</sup>	Substrate	Location	Collector	GenBank accession No. <sup>2</sup>			Reference
					tef1	his3	cmqA	
	CBS 112934; CMW 30987; CPC 4516	Soil under <i>Eucalyptus</i> forest	Northern Sumatra, Indonesia	M.J. Wingfield	AY725735	AY725698	AY725773	AY725651
<i>Ca. tetraramosa</i>	<b>CBS 136635;</b> CMW 31474; CERC 1809; CPC 23489	<i>E. urophylla</i> × <i>E.</i> <i>grandis</i> seedling leaf in nursery	SuiXi, ZhanJiang, GuangDong, China	G. Zhao	KJ462898	KJ463244	KJ463128	KJ463011
	CBS 136637; CMW 31476; CERC 1811	<i>E. urophylla</i> × <i>E.</i> <i>grandis</i> seedling leaf in nursery	SuiXi, ZhanJiang, GuangDong, China	G. Zhao	KJ462899	KJ463245	KJ463129	KJ463012
	<b>CBS 136077;</b> CMW 31411; CERC 1746; CPC 23479	Soil in <i>Eucalyptus</i> plantation	FangChengGang, GuangXi, China	X. Zhou, G. Zhao & F. Han	KJ462900	KJ463246	N/A	KJ463013
	CBS 136093; CMW 35410; CERC 1901	Soil in <i>Eucalyptus</i> plantation	GuangXi, China	X. Mou & R. Chang	KJ462901	KJ463247	KJ463130	KJ463014
	<b>CBS 125268;</b> CMW 9188	<i>E. grandis</i> cutting	Kwa-Zulu Natal, South Africa	L. Lombard	FJ972483	FJ972433	GQ267459	FJ972414
	CBS 125272; CMW 9896	<i>E. urophylla</i> × <i>E.</i> <i>grandis</i> cutting	Kwa-Zulu Natal, South Africa	L. Lombard	FJ972484	FJ972434	GQ267460	FJ972415

<sup>1</sup> CBS: Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands; CERC: China Eucalypt Research Centre, Zhanjiang, GuangDong Province, China; CMW: culture collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa; CPC: Pedro Crous working collection housed at CBS; HGUP: Plant Pathology Herbarium of Gui Zhou University, GuiYang 550025, China; IMI: CABI-Bioscience, Egham, UK.

<sup>2</sup> *tef1* = translation elongation factor 1-alpha; *his3* = histone H3; *cmqA* = calmodulin; *tub2* = β-tubulin.

<sup>3</sup> Isolates represent ex-type and indicated in **bold**.

<sup>4</sup> "N/A" represents sequences that are not available.

<sup>5</sup> "N/A" represents collectors that are not available.

**Supplementary Table 3.**

Supplementary Table 3. Statistics resulting from phylogenetic analyses.

Dataset	Phylogenetic group	No. of taxa	No. of bp <sup>1</sup>	Maximum parsimony			
				PI <sup>C</sup> <sup>2</sup>	No. of trees	Three length	CI <sup>3</sup>
tef1/his3/cmdA/tub2	Prolate	51	1993	579	184	861	0.8513
tef1/his3/cmdA/	Sphaero-Naviculate	58	1415	400	1166	620	0.8113
							0.9750
							0.9644
							0.8301
							0.7824
							0.1487
							0.1887

Dataset	Phylogenetic group	Subst. model <sup>7</sup>	NST <sup>8</sup>	Maximum likelihood		Rates
				Rate matrix	Rate matrix	
tef1/his3/cmdA/tub2	Prolate	TrN+G	6	10.000	25.771	10.000
tef1/his3/cmdA/	Sphaero-Naviculate	TIM2+I+G	6	14.897	44.198	14.897
				10.000	10.000	10.000
				41.384	41.384	41.384
				72.666	72.666	72.666
				Gamma	Gamma	Gamma

1 bp = base pairs.

2 PIC = number of parsimony informative characters.

3 CI = consistency index.

4 RI = retention index.

5 RC = rescaled consistency index.

6 HI = homoplasy index.

7 Subst. model = best fit substitution model.

8 NST = number of substitution rate categories.

## Supplementary Tables 4, 5, 6, and 7

**Supplementary Table 4.** Number of unique alleles found in *Calonectria aciculata*, *Ca. honghensis* and their phylogenetically closest related species in total and in the four gene regions.

	<i>Ca. honghensis</i>	<i>Ca. colhounii</i>	<i>Ca. monticola</i>
<i>Ca. aciculata</i>	12(3/1/3/5) <sup>1</sup>	18(10/1/5/2)	23(12/NA <sup>2</sup> /7/4)
<i>Ca. honghensis</i>		14(9/0/2/3)	19(13/NA/4/2)
<i>Ca. colhounii</i>			26(17/NA/6/3)

<sup>1</sup> The order of the four genes: total (*tef1*, *his*, *cmdA* and *tub2*).

<sup>2</sup> "NA" represents sequences that are not available.

**Supplementary Table 5.** Number of unique alleles found in *Ca. pseudoyunnanensis*, *Ca. yunnanensis* and their phylogenetically closest related species in total and in the four gene regions.

	<i>Ca. yunnanensis</i>	<i>Ca. asiatica</i>	<i>Ca. colombiensis</i>
<i>Ca. pseudoyunnanensis</i>	7(0/3/0/4) <sup>1</sup>	11(2/4/0/5)	27(3/9/2/13)
<i>Ca. yunnanensis</i>		12(2/4/0/6)	28(3/9/2/14)
<i>Ca. asiatica</i>			18(1/7/2/8)

<sup>1</sup> The order of the four genes: total (*tef1*, *his*, *cmdA* and *tub2*).

**Supplementary Table 6.** Number of unique alleles found in *Ca. lantauensis* and its phylogenetically closest related species in total and in the four gene regions.

	<i>Ca. curvispora</i>	<i>Ca. ilicicola</i>	<i>Ca. pacifica</i>	<i>Ca. sumatrensis</i>
<i>Ca. lantauensis</i>	36(4/15/17/NA <sup>1</sup> ) <sup>2</sup>	15(3/11/1/NA)	16(5/10/1/NA)	26(4/20/2/NA)
<i>Ca. curvispora</i>		28(5/7/16/NA)	28(1/11/16/NA)	44(4/23/17/NA)
<i>Ca. ilicicola</i>			16(6/10/0/NA)	25(5/19/1/NA)
<i>Ca. pacifica</i>				26(5/20/1/NA)

<sup>1</sup> "NA" represents sequences that are not available.

<sup>2</sup> The order of the four genes: total (*tef1*, *his*, *cmdA* and *tub2*).

**Supplementary Table 7.** Number of unique alleles found in *Ca. pseudoturangicola* and its phylogenetically closest related species in total and in the four gene regions.

	<i>Ca. hongkongensis</i>	<i>Ca. turangicola</i>
<i>Ca. pseudoturangicola</i>	34(7/13/4/10) <sup>1</sup>	6(3/0/2/1)
<i>Ca. hongkongensis</i>		30(4/13/4/9)

<sup>1</sup> The order of the four genes: total (*tef1*, *his*, *cmdA* and *tub2*).