

## Focus on azithromycin

N Padayachee,<sup>1</sup> M Milne,<sup>2</sup> N Schellack<sup>3</sup>

<sup>1</sup> Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, South Africa

<sup>2</sup> School of Pharmacy, Sefako Makgatho Health Sciences University, South Africa

<sup>3</sup> Department of Pharmacology, University of Pretoria, South Africa

**Corresponding author, email:** [neelaveni.padayachee@wits.ac.za](mailto:neelaveni.padayachee@wits.ac.za)

Azithromycin is a second-generation macrolide that was developed in the early 1980s as a semi-synthetic derivative of erythromycin. This broad-spectrum antibiotic is known for its activity against some Gram-negative organisms such as *H. influenza*. Azithromycin is similar to other macrolides, however, it also has the ability to inhibit quorum sensing and the formation of biofilm. Azithromycin has been beneficial in the treatment of influenza and Middle East respiratory syndrome coronavirus (MERS-CoV) and recently has shown to be effective against severe acute respiratory syndrome coronavirus 2 (SARS CoV-2) when used in combination with hydroxychloroquine or chloroquine. The side-effect that needs to be monitored carefully is the risk of cardiotoxicity when azithromycin is combined with chloroquine or hydroxychloroquine.

**Keywords:** azithromycin, COVID-19, antibacterial, antiviral, immunomodulatory, chloroquine, hydroxychloroquine

### Azithromycin

Azithromycin (AZM) was synthesised in the 1980s as a semi-synthetic derivative of erythromycin. This second-generation macrolide gained popularity due to its increased effects on host-defences and chronic human diseases. Furthermore, its popularity increased as it showed improved spectrum of activity, tissue pharmacokinetic characteristics and stability in an acid environment compared to erythromycin. It is also known for its activity against some Gram-negative organisms such as *Haemophilus influenza*.<sup>1-3</sup> Although azithromycin has the same antibacterial mechanism of action as other macrolides, its ability to inhibit bacterial quorum sensing, formation of biofilm and mucus production further increased its esteem.<sup>1</sup>

### Chemistry

In order to provide AZM with acid stability, it consists of a 15-membered macrocyclic lactone ring, with a sugar and an amino sugar attached, compared to the 14-membered macrocyclic lactone ring of the other macrolides. The alcohol and tertiary amino groups on the amino sugar (desosamine) are crucial for activity.<sup>5</sup> One major disadvantage of AZM is the

fact that it is very poorly soluble in aqueous environments ( $\pm 0.1 \text{ mgml}^{-1}$ ), which contributes to its relatively low absolute oral bioavailability of only 37%.<sup>6-8</sup>


Poor water solubility could lead to variable dissolution rates, and ultimately it will detrimentally influence the bioavailability and subsequently the treatment effectiveness of patients.<sup>9</sup>

The amorphous solid-state form of AZM has shown improved aqueous solubility and intestinal membrane permeability. This improved solubility ranged from 1.4 to 4.2 times the solubility of the crystalline form compared to the amorphous solid-state form.<sup>9</sup>

### Pharmacology

An additional methyl-substituted nitrogen atom combined into the lactone ring of erythromycin makes AZM a 15-membered lactone ring.<sup>10</sup> Its main role is on protein synthesis whereby it acts by inhibiting bacterial protein synthesis through binding to the 50S ribosomal subunit of susceptible organisms.<sup>11</sup> AZM's convenient once-daily dosing for 3–5 days is due to its long half-life thus sustaining high concentration of the drug in the tissue.

A 500 mg single dose of AZM achieves 37% bioavailability while the drugs' delivery into tissues is up to 100-fold higher than in the plasma.<sup>1,12,13</sup> Furthermore, AZM's concentrations are high in macrophages and polymorphonuclear leucocytes, tonsils, lung, prostate, liver and lymph nodes while fat and muscle displayed lower concentrations.<sup>3</sup> These high tissue and intracellular concentration levels are dependent on the dosing schedule of AZM.<sup>3,13</sup> AZM has shown high concentrations in infected fluids and tissues even after there are reduced concentrations in the plasma.<sup>13</sup>



**Figure 1:** Molecular structure of AZM<sup>4</sup>

## Antimicrobial spectrum

Due to the similarity of AZM to erythromycin, the antimicrobial spectrum activity extends to erythromycin susceptible, Gram-positive organisms and is much more effective against Gram-negative bacteria compared to the other macrolides.<sup>11</sup>

## Clinical uses

AZM is effective in the treatment of a multitude of conditions.<sup>11</sup>

The benefits of AZM have shown clinical effectiveness against many respiratory viral conditions such as influenza and Middle East respiratory syndrome coronavirus (MERS-CoV) and thus its use against the severe acute respiratory syndrome coronavirus 2 (SARSCoV-2) that causes the coronavirus disease-19 (COVID-19) is of no surprise. Furthermore, *in vitro* activity of AZM against viruses such as Ebola, Zika, rhinovirus and influenza has also been shown.<sup>14</sup>

## Drug interactions

AZM has been considered to be a safe drug with adverse drug reactions that affect the gastrointestinal system and central and peripheral nervous system. Ventricular depolarisation (QTc) interval prolongation, torsade de pointes (TdP), ventricular tachycardia and sudden cardiac death have been associated with macrolides, however, the incidences have been low globally. Cardiotoxicity in patients with SARS-CoV-2 pneumonia, however, has been a concern. Furthermore, the combination use of hydroxychloroquine and AZM in hospitalised patients has shown a higher risk of cardiac adverse events with risk being more likely related to the hydroxychloroquine.<sup>14</sup> Unlike other macrolide antibiotics such as erythromycin and clarithromycin, AZM is only a weak cytochrome P450 inhibitor.<sup>14</sup>

## Contraindications<sup>14</sup>

- Hepatic dysfunction
- Cholestatic jaundice
- Sensitivity to macrolides

## Caution<sup>14</sup>

- Hepatotoxicity
- Infantile hypertrophic pyloric stenosis
- Clostridioides difficile-associated diarrhoea
- Myasthenia gravis (this antibiotic may exacerbate muscle weakness)
- Patients with the previous prolongation of QTc interval or TdP

## Azithromycin for COVID-19

In contrast to chloroquine (CQ) or hydroxychloroquine (HCQ), AZM antiviral activity has been shown *in vitro* and/or *in vivo* on a large panel of viruses: Ebola, Zika, respiratory syncytial virus, influenza H<sub>1</sub>N<sub>1</sub> virus, enterovirus, and rhinovirus.<sup>15-24</sup> Its activity against respiratory syncytial virus has been demonstrated in a randomised study in infants.<sup>21</sup> AZM exhibited a synergistic antiviral effect against SARS-CoV-2 when combined with HCQ both *in vitro*,<sup>22</sup> and in a clinical setting.<sup>24</sup> Andreani et al. also reported a significant antiviral effect of AZM alone on SARS-

CoV-2.<sup>22</sup> The mechanisms of the antiviral effect of AZM support a large-spectrum of antiviral activity. AZM appears to decrease the virus entry into cells.<sup>19,25</sup> In addition, it can enhance the immune response against viruses by several actions. AZM up-regulates the production of type I and III interferons (especially interferon- $\beta$  and interferon- $\lambda$ ), and genes involved in virus recognition such as melanoma differentiation-associated (MDA)5 gene and retinoic acid-inducible gene (RIG)-I.<sup>18,23,24,26,27</sup> These mechanisms are universally involved in the innate response against infectious agents, and potentially against SARS-CoV-2. The immunomodulation properties of AZM are the rationale of its use against inflammatory manifestations leading to interstitial lung disease.<sup>28,29</sup>

AZM regulates and/or decreases the production of interleukin (IL)-1 $\beta$ , IL-6, IL-8, IL-10, IL-12, and Interferon alfa (IFN- $\alpha$ ).<sup>21,30,31</sup> AZM and HCQ both decrease the production of major inflammatory cytokines such as IL-1 and IL-6. However, the different profiles of immunomodulation between the two drugs may be crucial for selecting one of them for the treatment of COVID-19, in relation to the pathogenicity of the virus. Indeed, HCQ may decrease IL-2 levels but not AZM, while AZM may decrease IL-8 levels but not HCQ. AZM could allow a sufficient memory T-cell count to be maintained and therefore better immunisation.<sup>32</sup> Another property of AZM is its antibacterial effect, which may be most interesting to prevent or treat co-infection by bacteria and SARS-CoV-2. Recent data suggested that anaerobic bacteria of lung microbiota may be involved in the SARS-CoV-2 pathogenesis. Prevotella cells, which have been found in abnormal quantities in patients with severe disease, could internalise SARS-Cov2 and enhance its pathogenicity.<sup>33-35</sup> AZM is a possible treatment for Prevotella infections and decreases Prevotella-induced inflammation.<sup>36,37</sup> In addition, AZM is extensively distributed into tissue, especially in the lungs.<sup>38</sup>

## Conclusion

Azithromycin acts as an antibacterial, antiviral and an immunomodulator. It is used successfully in the treatment of respiratory conditions, urethritis and recently in COVID-19. Although most of the side-effects are manageable, there is a cardiovascular risk especially when used in combination with HCQ or CQ and patients need to be monitored carefully.

## ORCID

N Padayachee  <https://orcid.org/0000-0002-6146-8702>

M Milne  <https://orcid.org/0000-0002-9982-029X>

N Schellack  <https://orcid.org/0000-0001-9690-6285>

## References

1. Parnham MJ, Haber VE, Giamarellos-Bourboulis EJ, et al. Azithromycin: mechanisms of action and their relevance for clinical applications. *Pharmacology & Therapeutics*. 2014;143(2):225-45. <https://doi.org/10.1016/j.pharmthera.2014.03.003>.
2. Dunn CJ, Barradell LB. Azithromycin. *Drugs*. 1996;51(3):483-505. <https://doi.org/10.2165/00003495-199651030-00013>.
3. Lode H, Borner K, Koeppel P, Schaberg T. Azithromycin—review of key chemical, pharmacokinetic and microbiological features. *Journal of Antimicrobial Chemotherapy*. 1996;37(Suppl C):1-8. [https://doi.org/10.1093/jac/37.suppl\\_c.1](https://doi.org/10.1093/jac/37.suppl_c.1).

4. Moreton RC. United States Pharmacopeia-National Formulary. Journal of Excipients and Food Chemicals. 2013 Oct 2;6(3):925. <https://doi.org/10.32388/2yh02t>.
5. Patrick GL. An introduction to medicinal chemistry. Oxford University Press; 2013 Jan 10. <https://doi.org/10.1021/jm960427+>.
6. Adrjanowicz K, Zakowiecki D, Kaminski K, et al. Molecular dynamics in supercooled liquid and glassy states of antibiotics: azithromycin, clarithromycin and roxithromycin studied by dielectric spectroscopy. Advantages given by the amorphous state. Molecular Pharmaceutics. 2012 Jun 4;9(6):1748-63. <https://doi.org/10.1021/mp300067r>.
7. Montejo-Bernardo J, Garcia-Granda S, Bayod-Jasanada M, Llorente I, Llavonab L. An easy and general method for quantifying Azithromycin dihydrate in a matrix of amorphous Azithromycin. Arkivoc. 2005 Jan 1;9:321-31. <https://doi.org/10.3998/ark.5550190.0006.927>.
8. Arora SC, Sharma PK, Irchhaiya R, et al. Development, characterization and solubility study of solid dispersions of azithromycin dihydrate by solvent evaporation method. Journal of Advanced Pharmaceutical Technology & Research. 2010 Apr;1(2):221. <https://doi.org/10.4103%2F0110-5558.72427>.
9. Aucamp M, Odendaal R, Liebenberg W, Hamman J. Amorphous azithromycin with improved aqueous solubility and intestinal membrane permeability. Drug Development and Industrial Pharmacy. 2015 Jul 3;41(7):1100-8. <https://doi.org/10.3109/03639045.2014.93196>.
10. Bakheit AH, Al-Hadiya BM, Abd-Elgalil AA. Azithromycin. Profiles of Drug Substances, Excipients and Related Methodology. 2014;39:1-40. <https://doi.org/10.1016/B978-0-12-800173-8.00001-5>.
11. Martin DH, Mroczkowski TF, Dalu ZA, et al. A controlled trial of a single dose of azithromycin for the treatment of chlamydial urethritis and cervicitis. New England Journal of Medicine. 1992;327(13):921-5. <https://doi.org/10.1056/NEJM199209243271304>.
12. Nahata M. Drug interactions with azithromycin and the macrolides: an overview. Journal of Antimicrobial Chemotherapy. 1996;37(Suppl C):133-42. [https://doi.org/10.1093/jac/37.suppl\\_C.133](https://doi.org/10.1093/jac/37.suppl_C.133).
13. Langtry HD, Balfour JA. Azithromycin. Drugs. 1998 Aug;56(2):273-97. <https://doi.org/10.2165/00003495-199856020-00014>.
14. Echeverria-Esnal D, Martin-Ontiyuelo C, Navarrete-Rouco ME, et al. Azithromycin in the treatment of COVID-19: a review. Expert Review of Anti-Infective Therapy. 2020;19(2):147-63. <https://doi.org/10.1080/14787210.2020.1813024>.
15. Du X, Zuo X, Meng F, et al. Combinatorial screening of a panel of FDA-approved drugs identifies several candidates with anti-Ebola activities. Biochemical and Biophysical Research Communications. 2020;522(4):862-8. <https://doi.org/10.1016/j.bbrc.2019.11.065>.
16. Retalack H, Di Lullo E, Arias C, et al. Zika virus cell tropism in the developing human brain and inhibition by azithromycin. Proceedings of the National Academy of Sciences. 2016;113(50):14408-13. <https://doi.org/10.1073/pnas.1618029113>.
17. Wu Y-H, Tseng C-K, Lin C-K, et al. ICR suckling mouse model of Zika virus infection for disease modeling and drug validation. PLoS Neglected Tropical Diseases. 2018;12(10):e0006848. <https://doi.org/10.1371/journal.pntd.0006848>.
18. Li C, Zu S, Deng Y-Q, et al. Azithromycin protects against Zika virus infection by upregulating virus-induced type I and III interferon responses. Antimicrobial Agents and Chemotherapy. 2019;63(12):e00394-19. <https://doi.org/10.1128/aac.00394-19>.
19. Tran DH, Sugamata R, Hirose T, et al. Azithromycin, a 15-membered macrolide antibiotic, inhibits influenza A(H1N1)pdm09 virus infection by interfering with virus internalization process. The Journal of Antibiotics. 2019;72(10):759-68. <https://doi.org/10.1038/s41429-019-0204-x>.
20. Mosquera RA, De Jesus-Rojas W, Stark JM, et al. Role of prophylactic azithromycin to reduce airway inflammation and mortality in a RSV mouse infection model. Pediatric Pulmonology. 2018;53(5):567-74. <https://doi.org/10.1002/ppul.23956>.
21. Beigelman A, Isaacson-Schmid M, Sajol G, et al. Randomized trial to evaluate azithromycin's effects on serum and upper airway IL-8 levels and recurrent wheezing in infants with respiratory syncytial virus bronchiolitis. Journal of Allergy and Clinical Immunology. 2015;135(5):1171-8. <https://doi.org/10.1016/j.jaci.2014.10.001>.
22. Andreani J, Le Bideau M, Duflot I, et al. In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect. Microbial Pathogenesis. 2020;145:104228. <https://doi.org/10.1016/j.micpath.2020.104228>.
23. Zeng S, Meng X, Huang Q, et al. Spiramycin and azithromycin, safe for administration to children, exert antiviral activity against enterovirus A71 in vitro and in vivo. International Journal of Antimicrobial Agents. 2019;53(4):362-9. <https://doi.org/10.1016/j.ijantimicag.2018.12.009>.
24. Gautret P, Lagier J-C, Parola P, et al. Clinical and microbiological effect of a combination of hydroxychloroquine and azithromycin in 80 COVID-19 patients with at least a six-day follow up: a pilot observational study. Travel Medicine and Infectious Disease. 2020;34:101663. <https://doi.org/10.1016/j.tmaid.2020.101663>.
25. Yao X, Ye F, Zhang M, et al. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Clinical Infectious Diseases. 2020;71(15):732-9. <https://doi.org/10.1093/cid/ciaa237>.
26. Schögl A, Kopf BS, Edwards MR, et al. Novel antiviral properties of azithromycin in cystic fibrosis airway epithelial cells. European Respiratory Journal. 2015;45(2):428-39. <https://doi.org/10.1183/09031936.00102014>.
27. Menzel M, Akbarshahi H, Bjermer L, Uller L. Azithromycin induces anti-viral effects in cultured bronchial epithelial cells from COPD patients. Scientific Reports. 2016;6(1):28698. <https://doi.org/10.1038/srep28698>.
28. Kawamura K, Ichikado K, Suga M, Yoshioka M. Efficacy of azithromycin for treatment of acute exacerbation of chronic fibrosing interstitial pneumonia: a prospective, open-label study with historical controls. Respiration. 2014;87(6):478-84. <https://doi.org/10.1159/000358443>.
29. Bush A, Cunningham S, De Blic J, et al. European protocols for the diagnosis and initial treatment of interstitial lung disease in children. Thorax. 2015 Nov 1;70(11):1078-84. <https://doi.org/10.1136/thoraxjnl-2015-207349>.
30. Cai M, Bonella F, Dai H, et al. Macrolides inhibit cytokine production by alveolar macrophages in bronchiolitis obliterans organizing pneumonia. Immunobiology. 2013;218(6):930-7. <https://doi.org/10.1016/j.imbio.2012.10.014>.
31. Zargoulidis P, Papanas N, Kioumis I, et al. Macrolides: from in vitro anti-inflammatory and immunomodulatory properties to clinical practice in respiratory diseases. European Journal of Clinical Pharmacology. 2012;68(5):479-503. <https://doi.org/10.1007/s00228-011-1161-x>.
32. Tomazic J, Kotnik V, Wraber B. In vivo administration of azithromycin affects lymphocyte activity in vitro. Antimicrobial Agents and Chemotherapy. 1993;37(9):1786-9. <https://doi.org/10.1128/aac.37.9.1786>.
33. Chakraborty YS. Sequencing data (N = 3) shows Wuhan coronavirus integration in bacteria (Prevotella mostly). Sequencing arti-fact—or is the virus infecting both bacterial and human cells?. OSFPreprint. 2020. <https://doi.org/10.31219/osf.io/ktngw>.
34. Chakraborty S. The Wuhan coronavirus has integrated in Prevotella, which possibly causes the observed extreme virulence—as sequencing data from 2 different studies in China and Hong-Kong shows unequivocally. OSFPreprint. 2020. <https://doi.org/10.31219/osf.io/ktngw>.
35. Chakraborty S. The 2019 Wuhan outbreak could be caused by the bacteria Prevotella, which is aided by the coronavirus, possibly to adhere to epithelial cells – Prevotella is present in huge amounts in patients from both China and Hong Kong. <https://doi.org/10.4103/1755-6783.190196>.
36. Mättö J, Asikainen S, Väisänen ML, et al.  $\beta$ -lactamase production in Prevotella intermedia, Prevotella nigrescens, and Prevotella pallens. Genotypes and in vitro susceptibilities to selected antimicrobial agents. Antimicrobial Agents and Chemotherapy. 1999;43(10):2383-8. <https://doi.org/10.1128/AAC.43.10.2383>.
37. Choi E-Y, Jin J-Y, Choi J-I, Choi IS, Kim S-J. Effect of azithromycin on Prevotella intermedia lipopolysaccharide-induced production of interleukin-6 in murine macrophages. European Journal of Pharmacology. 2014;729:10-6. <https://doi.org/10.1016/j.ejphar.2014.02.008>.
38. Rodvold KA, Gotfried MH, Danziger LH, Servi RJ. Intrapulmonary steady-state concentrations of clarithromycin and azithromycin in healthy adult volunteers. Antimicrobial Agents and Chemotherapy. 1997;41(6):1399-402. <https://doi.org/10.1128/aac.41.6.1399>.