PTSD: PC-PTSD-5 astance Use Screening Alcohol: AUDIT-C Alcohol: CAGE	Course Modules	Question Clir Bank Cha	hical Tools & Calculators >	Clinical Consultation	HIV Resources >
 Permentia: HDS This tool is gg meant to be viewed by the patient during the test. The provider should use it as a guide and all actions to the patient with their own hands. The timers will entit adult out case at the beginning and end of the time period. To perform this test follow the steps in order below: Memory-Registration Memory-Registration I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. Then as the patient will off adult of the patient dues on target adult the patient will off adult of the patient will off adult off. I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. The near store patient will off adult off. I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. The near store patient will off adult off. I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. The near store patient will off adult off. I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. The near store patient words after you have said them. Reperent of the patient tag the first two fingers of the non-dominant hand as widely and as quickly as possible for the patient tag the first two fingers of the non-dominant hand as widely and as quickly as possible for the calculator The calculator The calculator Gue table patient of taps performed in 5 seconds: Seconds 					Share
anxiety: GAD-2 Anxiety: GAD-7 Depression: PHQ-2 Depression: PHQ-9 PTSD: PC-PTSD-5 stance Use Screening Akohol: AUDIT-C Akohol: AUDIT-C Akohol: CAGE CAGE-AID Drug Abuse: TICS Dipuid: Risk Tool Dica Abuse: TICS Dipuid: Risk Tool Dica Calculator SMC CALCULAT	DS	5			use it as a guide and demonstrat
Anviety: GAD-7 Depression: PHQ-2 Depression: PHQ-3 Perform this test follow the steps in order below: Memory-Registration Item ask the patient does not recall (dog, hat, bean, red) - 1 second to say each. 1. Give four words to recall (dog, hat, bean, red) - 1 second to say each. 2. Then ask the patient does not recall them all immediately. 3. Repeat words if the patient does not recall them all immediately. 4. Tell the patient does not recall of the words again a bit later. Nethol: CAGE Auchol: CAGE Auchol: CAGE Auchol: CAGE Auchol: CAGE Auchol: CAGE Auchol: CAGE Drug Abuse: TICS Diploid: Risk Tool Iteal Calculator RPRI Calculator TP Calculator Count the number of taps performed in 5 seconds:	ā	all actions to the patient wi	th their own hands.	·	
dog hat bean rec PrSD: PC-PTSD-5 I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. I. Then ask the patient all four words after you have said them, Bepression: PHQ-9 I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. I. Then ask the patient all four words after you have said them, Bepression: PHQ-9 I. Give four words to recall (dog, hat, bean, red) - 1 second to say each. I. Then ask the patient all four words after you have said them, Bepression: PHQ-9 I. Have the patient you will ask for recall of the words again a bit later. I. Motor Speed O-4 p Have the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilitical Calculator Seconds Index of taps performed in 5 seconds: Seconds				inte period.	
dog nat Dean red prsD; PC-PTSD-5 .	IQ-2	emory-Registration			
 TSD: PC-PTSD-5 stance Use Screening Neohol: AUDIT-C Neohol: AUDIT-C Neohol: CAGE Then ask the patient all four words after you have said them. Repeat words if the patient does not recall them all immediately. Tell the patient you will ask for recall of the words again a bit later. Notor Speed O-4 program a suidely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possibilities in the patient tap the first two fingers of the non-dominant hand as widely and as quickly as po	IQ-9	dog	hat	bean	red
stance Use Screening Alcohol: AUDIT-C Alcohol: AUDIT-C Alcohol: AUDIT-C Alcohol: AUDIT-C Alcohol: CAGE CAGE-AID Drug Abuse: DAST-10 Drug Abuse: TICS Dirug Abuse: TICS Secouldator TIP Calculator TIP Calculator SFR Calculator SFR Calculator	-5				
4. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A. Tell the patient you will ask for recall of the words again a bit later. A.	creening	2. Then ask the patient all t	four words after you have said them,	-	
TAGE-AID 0-4 pr Drug Abuse: DAST-10 1. Motor Speed Drug Abuse: TICS 0-1 pr Dpioid: Risk Tool Image: Comparison of the non-dominant hand as widely and as quickly as possible ical Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible MPRI Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible TrC Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible TrC Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible TrC Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible TrC Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible Image: Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible Image: Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible Image: Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible Image: Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible Image: Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible Image: Calculator Image: Comparison of the non-dominant hand as widely and as quickly as possible Image: Calculator Image: Comparison of the non-dominant hand as widely and the non-do	- C	4. Tell the patient you will a	ask for recall of the words again a bit	later.	
Have the patient tap the first two fingers of the non-dominant hand as widely and as quickly as possible Drug Abuse: DAST-10 Drug Abuse: TICS Dpioid: Risk Tool ical Calculators APRI Calculator TrCI Calculator TrCI Calculator TPO 4 Calculator TEPO4 Calculator SFR CALCULATOR S					0.4 points
Drug Abuse: TICS Dpioid: Risk Tool ical Calculators MRI Calculator SMI Calculator TTP Calculator TEPO4 Calculator GFR Calculator SFR Calculator	1.1	Motor Speed			0-4 points
Dpioid: Risk Tool ical Calculators APRI Calculator BMI Calculator CrCl Calculator TP Calculator TP Calculator FIB-4 Calculator GFR Calculator SFR Calculator SFR Calculator Count the number of taps performed in 5 seconds: 5 seconds					
ical Calculators APRI Calculator CrCl Calculator TP Calculator FEPO4 Calculator FEPO4 Calculator GFR Calculator FEPO4 CALCULA	AST-10 Ha	ve the patient tap the fir	st two fingers of the non-dominan	: hand as widely and	as quickly as possible.
APRI Calculator BMI Calculator CrCI Calculator FIB-4 Calculator FEPO4 Calculator GFR Calculator GFR Calculator Count the number of taps performed in 5 seconds: 5 seconds	AST-10	ve the patient tap the fir	st two fingers of the non-dominan	hand as widely and	as quickly as possible.
BMI Calculator CrCl Calculator CTP Calculator FIB-4 Calculator FEPO4 Calculator GFR Calculator GFR Calculator	CS	ve the patient tap the fir.	st two fingers of the non-dominan	hand as widely and	as quickly as possible.
CrCl Calculator CTP Calculator FIB-4 Calculator FEPO4 Calculator GFR Calculator 5 seconds	ol	ve the patient tap the fir.	st two fingers of the non-dominant	hand as widely and	as quickly as possible.
CTP Calculator FIB-4 Calculator EEPO4 Calculator GFR Calculator 5 seconds: 5 seconds	asi-10 CS ol ors	ve the patient tap the fir.	st two fingers of the non-dominant \leftarrow	hand as widely and	as quickly as possible.
FIB-4 Calculator Count the number of taps performed in 5 seconds: GFR Calculator	ASI-10 CS ol ors	ve the patient tap the fir.	st two fingers of the non-dominant \leftarrow	hand as widely and	as quickly as possible.
EPO4 Calculator Count the number of taps performed in 5 seconds: GFR Calculator 5 seconds	ASI-10 CS ol ors	ve the patient tap the fir.	st two fingers of the non-dominant	hand as widely and	as quickly as possible.
GFR Calculator 5 seconds	ASI-10 CS ol ors	ve the patient tap the fir.	st two fingers of the non-dominant \leftarrow	hand as widely and	as quickly as possible.
	ASI-10 CS ol ors	ve the patient tap the fir.	st two fingers of the non-dominant \leftarrow	hand as widely and	as quickly as possible.
	ASI-10 CS ol ors		\rightarrow	hand as widely and	as quickly as possible.
	ASI-10 CS ol ors r r r cor Co		ps performed in 5 seconds: 5 second	s	as quickly as possible.
Scoring: Record Number of Taps: ≥ 15 taps = 4 points 11-14 taps = 3 points	ASI-10 CS ol ors r r r cor Co		ps performed in 5 seconds: 5 second	s	2

- - •

· -

.

International HIV Dementia Scale (IHDS) - Mental Disorders Screening - National HIV Curriculum

ements with the non-dominar	nt hand as quickly as
	3
fist on flat surface.	
-	it.
e.	
) seconds:	
econds	
rt Timer	
2 sequences 1 sequences	s = 2 points e = 1 point
	0-4 points
bean	red
clue as follows:	
	• •
piece of cloth hat	iing?
	ning?
	rface with palm down. surface on the side of the 5th dig se. 9 seconds: t Timer Scor ≥ 4 sequences 2 sequences 1 sequences

Interpretation:

• This is the sum of the scores on items 1-3. The maximum possible score is 12 points. A patient with a score of \leq 10 should be evaluated further for possible dementia.

Acknowledgement

- The International HIV Dementia Scale (IHDS) was developed by Dr. Ned C. Sacktor, Professor of Neurology, Johns Hopkins
 Medicine.
- Reproduced with permission from Dr. Ned C. Sacktor.

Sources

 Sacktor NC, Wong M, Nakasujja N, et al. The International HIV Dementia Scale: a new rapid screening test for HIV dementia. AIDS. 2005;19:1367-74

① This calculator operates entirely from your device.

No input variables or data is transmitted between your computer and our servers.

Funded by a grant from the Health Resources and Services Administration (HRSA)

HIV Medications

Single-Tablet Regimens Entry Inhibitors Nucleoside Reverse Transcriptase Inhibitors Non-Nucleoside Reverse Transcriptase Inhibitors Integrase Inhibitors Protease Inhibitors Pharmacokinetic Enhancers

Copyright © 2019 National HIV Curriculum Version 2.7.2-c - AWS

Course Modules

Screening and Diagnosis Basic HIV Primary Care Antiretroviral Therapy Co-Occurring Conditions Prevention of HIV Key Populations Question Bank Clinical Challenges Master Bibliography

Contributors Content Bundles

CME provided by

Tools & Calculators

Mental Disorders Screening Substance Use Screening Clinical Calculators

Contact Us | Terms and Conditions | Privacy Policy CE (CNE/CME) Notices | Copyright and Attribution Notices

This project is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number U10HA32104. No percentage of this project was financed with non-governmental sources. This information or content and conclusions are those of the authors and should not be construed as the official position or policy of, nor should any endorsements by inferred by HRSA, HHS or the U.S. Government.

part of the IDEA platform