

Supplementary Material

Spatial constraints and seasonal conditions but not poaching pressure are linked with elevated faecal glucocorticoid metabolite concentrations in white rhino

Zoliswa N. Nhleko^{A,B,}, Andre Ganswindt^C, Sam M. Ferreira^B, and Robert A. McCleery^D*

^AInterdisciplinary Program in Ecology, University of Florida, Gainesville, FL 32611, USA.

^BSavanna Node, Scientific Services, SANParks, Skukuza 1350, South Africa.

^CMammal Research Institute, University of Pretoria, Pretoria 0028, South Africa.

^DDepartment of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL 32611, USA.

*Correspondence to: Zoliswa N. Nhleko Interdisciplinary Program in Ecology, University of Florida, Gainesville, FL 32611, USA Email: nhlekozn@gmail.com

Appendix 1 Changes in fecal glucocorticoid metabolite (fGCM) concentrations post-defecation for samples stored at ambient temperature for up to 7 days. Dots at a given point in time represent the mean percentage change for the simultaneously frozen sample triplicates. The gray dashed line is the baseline level for the samples frozen at time zero ($\bar{x} = 0.54 \mu\text{g/g}$ dry weight).

